

## Notice of Weed Control Application

**Date of Application:** March 13-17, 2023

**Locations:**

- Flamingo Rd.
- Twain Ave.
- Desert Inn Rd.
- Hualapai Way.
- Navajo Willow Ln.
- Desert Primrose Ln.
- Spotted Leaf Ln.

**Reason for Application:** Target weed control in planters, rock areas, tree wells, and cracks of sidewalks.

**Product Manufacturer Name:** Ranger Pro Herbicide.

-EPA registration no. 524-517

-Active ingredients: glyphosate

-Signal word: Caution

-Precautionary statement: Harmful if swallowed or inhaled. Causes eye irritation.

**Product Manufacturer Name:** Pendulum Aqua Cap

-EPA registration no. 241-416

-Active ingredients: pendimethalin: N-(1ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine

-Signal word: Caution

-Precautionary statement: Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing.

## ATTENTION:

This specimen label is provided for general information only.

- This pesticide product may not yet be available or approved for sale or use in your area.
- It is your responsibility to follow all Federal, state and local laws and regulations regarding the use of pesticides.
- Before using any pesticide, be sure the intended use is approved in your state or locality.
- Your state or locality may require additional precautions and instructions for use of this product that are not included here.
- Monsanto does not guarantee the completeness or accuracy of this specimen label. The information found in this label may differ from the information found on the product label. You must have the EPA approved labeling with you at the time of use and must read and follow all label directions.
- You should not base any use of a similar product on the precautions, instructions for use or other information you find here.
- Always follow the precautions and instructions for use on the label of the pesticide you are using.

21225G1-13



## Complete Directions for Use

The complete broad-spectrum postemergence professional herbicide for industrial, turf and ornamental weed control.

EPA Reg. No. 524-517

2007-1

**AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.**

Read the entire label before using this product.

Use only according to label instructions.

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

## 1.0 INGREDIENTS

### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl)glycine,  
in the form of its isopropylamine salt..... 41.0%  
OTHER INGREDIENTS (including surfactant): ..... 59.0%  
100.0%

\*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent Nos. 5,683,958; 5,703,015; 6,063,733; 6,121,199; 6,121,200. No license granted under any non-U.S. patent(s).

## 2.0 IMPORTANT PHONE NUMBERS

FOR PRODUCT INFORMATION OR ASSISTANCE  
IN USING THIS PRODUCT,  
CALL TOLL-FREE, 1-800-332-3111.

IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT,  
OR FOR MEDICAL ASSISTANCE,  
CALL COLLECT, DAY OR NIGHT, (314)-694-4000.

## 3.0 PRECAUTIONARY STATEMENTS

## 3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children.

### CAUTION!

CAUSES EYE IRRITATION.

Avoid contact with eyes or clothing.

**FIRST AID:** Call a poison control center or doctor for treatment advice.

<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses if present after the first 5 minutes then continue rinsing eye.</li></ul>
-------------------	---

- Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
- You may also contact (314) 694-4000, collect day or night, for emergency medical treatment information.
- This product is identified as **Ranger PRO® herbicide, EPA Registration No. 524-517.**

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## 3.2 Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

## 3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Monsanto Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, chemical resistant gloves greater than 14 mils in thickness composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks.

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (40 CFR Part 170) for agricultural pesticides. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

## 4.0 STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

Keep container closed to prevent spills and contamination.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures.

**CONTAINER DISPOSAL:** Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

**FOR REFILLABLE PORTABLE CONTAINERS:** Do not reuse this container except for refill in accordance with a valid Monsanto Repackaging or Toll Repackaging Agreement. If not refilled or returned to the authorized repackaging facility, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FOR BULK CONTAINERS:** Triple rinse emptied bulk container. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

**FOR PLASTIC 1-WAY CONTAINERS & BOTTLES:** Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FOR DRUMS:** Do not reuse container. Return container per the Monsanto container return program. If not returned, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## 5.0 GENERAL INFORMATION (How This Product Works)

**Product Description:** This product is a postemergence, systemic herbicide with no soil residual activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid containing surfactant.

**Time to Symptoms:** This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

**Mode of Action in Plants:** The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

**Cultural Considerations:** Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the recommended stage for treatment.

**Rainfastness:** Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

**No Soil Activity:** Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

**Tank Mixing:** This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this label. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

When this label recommends a tank mixture with a generic active ingredient such as diuron, atrazine, 2,4-D, or dicamba, the user is responsible for ensuring that the mixture product's label allows the specific application.

**Annual Maximum Use Rate:** The combined total of all treatments must not exceed 10.6 quarts of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate containing products does not exceed stated maximum use rates.

### ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

**NOTE:** Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

## 6.0 MIXING

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

**NOTE:** REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS VISIBLY MUDDY WATER OR WATER FROM PONDS AND DITCHES THAT IS NOT CLEAR.

### 6.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of this product near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

### 6.2 Tank Mixing Procedure

When tank mixing, read and carefully observe label directions, cautionary statements and all information on the labels of all products used. Add the tank-mix product to the tank as directed by the label. Maintain agitation and add the recommended amount of this product.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation may be required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance. Ensure that the specific tank mixture product is registered for application at the desired site.

Refer to the "TANK MIXING" section of "GENERAL INFORMATION" for additional precautions.

## 6.3 Mixing for Hand-Held Sprayers

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

### Spray Solution

Desired Volume	Amount of Ranger PRO herbicide					
	1/2%	1%	1-1/2%	2%	5%	10%
1 gal	2/3 oz	1-1/3 oz	2 oz	2-2/3 oz	6-1/2 oz	13 oz
25 gal	1 pt	1 qt	1-1/2 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1-1/2 gal	2 gal	5 gal	10 gal

2 tablespoons = 1 fluid ounce

For use in backpack, knapsack or pump-up sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

## 6.4 Surfactant

Nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When adding additional surfactant, use 0.5-percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1-percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

## 6.5 Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilution. Use colorants or dyes according to the manufacturer's recommendations.

## 6.6 Drift Control Additives

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

# 7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

### SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

### AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

### Importance of droplet size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the **Wind**, **Temperature and Humidity**, and **Temperature Inversion** sections of this label).

### Controlling droplet size

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.

- **Pressure:** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

### Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller droplets, etc.).

### Wind

Drift potential is lowest between wind speeds of 2 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 miles per hour due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

### Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### Sensitive Areas

This product should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

## 7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

This product plus dicamba tank mixtures may not be applied by air in California.

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Avoid direct application to any body of water.

Use the recommended rates of this herbicide in 3 to 25 gallons of water per acre.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application—To avoid streaked, uneven or overlapped application, use appropriate marking devices.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion. To prevent corrosion of exposed parts, thoroughly wash aircraft after each day of spraying to remove residues of this product accumulated during spraying or from spills. Landing gear is most susceptible.

## 7.2 Ground Broadcast Equipment

Use the recommended rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check for even distribution of spray droplets.

## 7.3 Hand-Held or High-Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

For control of weeds listed in the "ANNUAL WEEDS" section of "WEEDS CONTROLLED", apply a 1/2-percent solution of this product to weeds less than 6 inches in height or runner length. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1-percent solution. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds.

For best results, use a 2-percent solution on harder-to-control perennials, such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

For low volume directed spray applications, use a 5- to 10-percent solution of this product for control or partial control of annual weeds, perennial weeds, or woody brush and trees. Spray coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

## 7.4 Selective Equipment

This product may be applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars, after dilution and thorough mixing with water, to listed weeds growing in any non-crop site specified on this label.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

**AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION, AS SERIOUS INJURY OR DEATH IS LIKELY TO OCCUR.**

Applicators used above desired vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation is likely to result in discoloration, stunting or destruction.

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

### Shielded and Hooded Applicators

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide. Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

### Wiper Applicators and Sponge Bars

A wiper or sponge applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

**For Rope or Sponge Wick Applicators**—Solutions ranging from 33 to 75 percent of this product in water may be used.

**For Panel Applicators and pressure-feed systems**—Solutions ranging from 33 to 100 percent of this product in water may be used.

When applied as recommended above, this product **CONTROLS** the following weeds:

Corn, volunteer	Sicklepod
Panicum, Texas	Spanishneedles
Rye, common	Starbur, bristly
Shattercane	

When applied as recommended above, this product **SUPPRESSES** the following weeds:

Beggarweed, Florida	Ragweed, common
Bermudagrass	Ragweed, giant
Dogbane, hemp	Smutgrass
Dogfennel	Sunflower
Guineagrass	Thistle, Canada
Johnsongrass	Thistle, musk
Milkweed	Vaseygrass
Nightshade, silverleaf	Velvetleaf
Pigweed, redroot	

## 7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the undiluted concentrate of other products when using injection systems unless specifically recommended.

## 7.6 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

CDA equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

## 8.0 SITE AND USE RECOMMENDATIONS

Detailed instructions follow alphabetically, by site.

Unless otherwise specified, applications may be made to control any weeds listed in the annual, perennial and woody brush tables. Refer also to the "SELECTIVE EQUIPMENT" section.

### 8.1 Cut Stumps

Cut stump treatments may be made on any site listed on this label. This product will control many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50- to 100-percent solution of this product to the freshly-cut surface **immediately after cutting**. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

Alder	Saltcedar
Eucalyptus	Sweetgum
Madrone	Tan oak
Oak	Willow
Reed, giant	

**DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP.** Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

### 8.2 Forestry Site Preparation

This product is recommended for the control or partial control of woody brush, trees and herbaceous weeds in forestry. This product is also recommended for use in preparing or establishing wildlife openings within these sites and maintaining logging roads.

This product is recommended for use in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

#### APPLICATION RATES AND TIMING

APPLICATION	RANGER PRO HERBICIDE	SPRAY VOLUME GAL/A
<b>BROADCAST</b>		
Aerial	2 to 10 qts/A	5 to 30
Ground	2 to 10 qts/A	10 to 60
<b>SPRAY-TO-WET</b>		
Handgun, Backpack	3/4% to 2%	spray-to-wet by volume
<b>LOW VOLUME DIRECTED SPRAY</b>		
Handgun, Backpack	5% to 10%	partial coverage* by volume

\*For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results.

Use higher rates of this product within the recommended range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the recommended range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the recommended range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

#### Tank Mixtures

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

**NOTE:** For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any recommended rate of this product may be used in a tank mix with the following products for forestry site preparation.

#### PRODUCT

Arsenal Applicators Concentrate

Escort

Chopper

Garlon 4

Oust

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release.

## 8.3 General Non-crop Areas and Industrial Sites

Use in areas such as airports, apartment complexes, ditch banks, dry ditches, dry canals, fencerows, golf courses, industrial sites, lumber yards, manufacturing sites, office complexes, parks, parking areas, petroleum tank farms and pumping installations, railroads, recreational areas, residential areas, roadsides, sod or turf seed farms, schools, storage areas, substations, warehouse areas, other public areas, and similar industrial and non-crop sites.

#### General Weed Control, Trim-and-Edge, Bare Ground

This product may be used in general non-crop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product may be tank mixed with the following products provided that the specific product is registered for use on such non-crop sites. Refer to these products' labels for approved non-crop sites and application rates.

Arsenal	Karmex DF	Ronstar 50 WSP
Barricade 65WG	Krovar I DF	Sahara
Clarity	Oust	simazine
diuron	Pendulum 3.3 EC	Surflan
Endurance	Pendulum WDG	Telar
Escort	Plateau	Vanquish
Garlon 3A	Princep DF	2,4-D
Garlon 4	Princep Liquid	

When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control or partial control of emerged perennial weeds, woody brush and trees.

For control or partial control of the following perennial weeds, apply 1 to 2 quarts of this product plus 2 to 4 ounces of Oust per acre.

Bahiagrass	Fescue, tall
Bermudagrass	Johnsongrass
Broomsedge	Poorjoe
Dallisgrass	Quackgrass
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

#### Chemical Mowing - Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 6 fluid ounces of this

product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

#### Chemical Mowing - Annuals

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 4 to 5 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

#### Bromus Species and Medusahead in Pastures and Rangelands

*Bromus* species. This product may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial, rangeland and pasture sites. Apply 8 to 16 fluid ounces of this product per acre on a broadcast basis.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses can become reestablished on the site.

Medusahead. To treat medusahead, apply 16 fluid ounces of this product per acre as soon as plants are actively growing, and prior to the 4-leaf stage. Applications may be made in the fall or spring.

Applications to brome and medusahead may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre. When applied as directed in this label, there are no grazing restrictions.

#### Dormant Turfgrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant Bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring green-up.

Apply 8 to 64 fluid ounces of this product per acre. Apply the recommended rates in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 16 fluid ounces per acre may result in injury or delayed green-up in highly maintained areas, such as golf courses and lawns. DO NOT apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for dormant Bermudagrass and bahiagrass treatments.

#### Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. DO NOT apply more than 16 fluid ounces of this product per acre in highly maintained turfgrass areas. DO NOT apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for actively growing Bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

#### Turfgrass Renovation, Seed, or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as Bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow translocation into underground plant parts.

Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

If application rates total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

## 8.4 Habitat Management

#### Habitat Restoration and Management

This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

### Wildlife Food Plots

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

## 8.5 Injection and Frill (Woody Brush and Trees)

This product may be used to control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 1/25 fluid ounce (1 mL) of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100 percent concentration of this product. For best results, application should be made during periods of active growth and after full leaf expansion. This product will control many species, some of which are listed below:

<u>Control</u>	<u>Partial Control</u>
Oak	Black gum
Poplar	Dogwood
Sweetgum	Hickory
Sycamore	Maple, red

## 8.6 Ornamentals, Plant Nurseries, and Christmas Trees

### Post-Directed, Trim-and-Edge

This product may be used as a post-directed spray around established woody ornamental species such as arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, douglas fir, joboba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew. This product may also be used to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a nursery setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material. **THIS PRODUCT IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES.** Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established ornamental species.

### Site Preparation

This product may be used prior to planting any ornamental, nursery or Christmas tree species.

### Wiper Applications

This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established eucalyptus or poplar trees. See the "SELECTIVE EQUIPMENT" section of this label for further information about the proper use of wiper applicators.

### Greenhouse/Shadehouse

This product may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

## 8.7 Parks, Recreational and Residential Areas

This product may be used in parks, recreational and residential areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around trees, fences, and paths, around buildings, sidewalks, and other objects in these areas. This product may be used for spot treatment of unwanted vegetation. This product may be used to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

All of the instructions in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section apply to park and recreational areas.

## 8.8 Railroads

All of the instructions in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section apply to railroads.

### Bare Ground, Ballast and Shoulders, Crossings, and Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight

at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.

This product may be tank mixed with the following products provided that the specific product is registered for ballast, shoulder, spot, bare ground and crossing treatments:

Arsenal	Krovar I DF
Clarity	Oust
diuron	Sahara
Escort	Spike
Garlon 3A	Telar
Garlon 4	Vanquish
Hyvar X	2,4-D

### Brush Control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 4 to 10 quarts of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 3/4- to 2-percent solution of this product when using high-volume spray-to-wet applications. Apply a 5- to 10-percent solution of this product when using low volume directed sprays for spot treatment. This product may be mixed with the following products for enhanced control of woody brush and trees:

Arsenal	Garlon 4
Escort	Tordon K
Garlon 3A	

### Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpet creeper
Fescue, tall	Vaseygrass

This product may be tank-mixed with Oust. If tank-mixed, use no more than 1 to 3 pints of this product with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Blackberry	Johnsongrass
Bluestem, silver	Poorjoe
Broomsedge	Raspberry
Dallisgrass	Trumpet creeper
Dewberry	Vaseygrass
Dock, curly	Vervain, blue
Dogfennel	

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may occur.

## 8.9 Roadsides

All of the instructions in the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section apply to roadsides.

### Shoulder Treatments

This product may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

### Guardrails and Other Obstacles to Mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

### Spot Treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

### Tank Mixtures

This product may be tank-mixed with the following products provided that the specific product is registered for shoulder, guardrail, spot and bare ground treatments:

Clarity	Princep DF
diuron	Princep Liquid
Endurance	Ronstar 50 WSP
Escort	Sahara
Krovar I DF	simazine
Oust	Surflan
Outrider®	Telar
Pendulum 3.3 EC	Vanquish
Pendulum WDG	2,4-D

See the "GENERAL NON-CROP AREAS AND INDUSTRIAL SITES" section of this label for general instructions for tank mixing.

#### Release of Bermudagrass or Bahiagrass

##### Dormant Applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant Bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring green-up. This product may also be tank-mixed with Outrider herbicide or Oust for residual control. Tank mixtures of this product with Oust may delay green-up.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

Apply 8 to 64 fluid ounces of this product in a tank mixture with 3/4 to 1 1/3 ounces Outrider herbicide per acre. Read and follow all label directions for Outrider herbicide.

Apply 8 to 64 fluid ounces of this product per acre alone or in a tank mixture with 1/4 to 1 ounce per acre of Oust. Apply the recommended rates in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in green-up and minimize injury, add no more than 1 ounce of Oust per acre on Bermudagrass and no more than 1/2 ounce of Oust per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

##### Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in 10 to 40 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpet creeper
Fescue, tall	Vaseygrass

This product may be tank mixed with Outrider herbicide for control or partial control of Johnsongrass and other weeds listed in the Outrider herbicide label. Use 8 to 32 fluid ounces of this product with 3/4 to 1 1/3 ounces of Outrider herbicide. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height.

This product may be tank-mixed with Oust. If tank-mixed, use no more than 1 to 2 pints of this product with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Bluestem, silver	Johnsongrass
Broomsedge	Poorjoe
Dallisgrass	Trumpet creeper
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not recommended, since severe injury may occur.

##### Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

This product may be used for control or partial control of Johnsongrass and other weeds listed on the Outrider herbicide label in actively growing bahiagrass. Apply 1 1/2 to 4 3/4 ounces of this product with 3/4 to 1 1/3 ounces of Outrider herbicide per acre. Use the higher rates for control of perennial weeds or annual weeds greater than 6 inches in height. Use only on well established bahiagrass.

A tank mixture of this product plus Oust may be used. Apply 6 fluid ounces of this product plus 1/4 ounce of Oust per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

## 8.10 Utility Sites

In utilities, this product is recommended for use along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, such as substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities.

This product is also recommended for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

#### Tank Mixtures

Tank mixtures of this product may be used to increase the spectrum of control for herbaceous weeds, woody brush and trees provided that the specific product is registered for application to the desired site. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any recommended rate of this product may be used in a tank mix.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

**NOTE:** For side trimming treatments, it is recommended that this product be used alone or in tank mixture with Garlon 4.

#### BROADCAST APPLICATION

Arsenal  
Escort  
Garlon 3A\*  
Garlon 4  
Oust

#### SPRAY-TO-WET APPLICATION

Arsenal  
Escort

#### LOW VOLUME DIRECTED SPRAY APPLICATION

Arsenal  
Escort

\*Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray compatibility problems.

#### Bare Ground and Trim-and-Edge

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product may be tank mixed with the following products. Refer to these products' labels for approved non-crop sites and application rates.

Arsenal	Plateau
Banvel	Princep DF
Barricade 65WG	Princep Liquid
diuron	Ronstar 50 WSP
Endurance	Sahara
Escort	simazine
Garlon 3A	Surflan

## 9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area.

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the following label sections for recommended rates for the control of annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 5 to 10 quarts per acre for enhanced results.

## 9.1 Annual Weeds

Use 1 quart per acre if weeds are less than 6 inches in height or runner length and 1.5 quarts to 4 quarts per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions.

For spray-to-wet applications, apply a 1/2-percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 1- to 2-percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall.

#### WEED SPECIES

Anoda, spurred	Brome, downy*
Barley*	Brome, Japanese*
Barnyardgrass*	Browntop panicum*
Bassia, fivehook	Buttercup*
Bittercress*	Carolina foxtail*
Black nightshade*	Carolina geranium
Bluegrass, annual*	Castorbean
Bluegrass, bulbous*	Cheatgrass*

## WEED SPECIES (Cont'd.)

Cheeseweed ( <i>Malva parviflora</i> )	Mustard, tansy*
Chervil*	Mustard, tumble*
Chickweed*	Mustard, wild*
Cocklebur*	Oats
Copperleaf, hophornbeam	Pigweed*
Corn*	Plains/Tickseed coreopsis*
Corn speedwell*	Prickly lettuce*
Crabgrass*	Puncturevine
Dwarf dandelion*	Purslane, common
Eastern manna grass*	Ragweed, common*
Eclipta*	Ragweed, giant
Fall panicum*	Red rice
Falsedandelion*	Russian thistle
Falseflax, smallseed*	Rye*
Fiddleneck	Ryegrass*
Field pennycress*	Sandbur, field*
Filaree	Shattercane*
Fleabane, annual*	Shepherd's-purse*
Fleabane, hairy	Sicklepod
( <i>Conyza bonariensis</i> )*	Signalgrass, broadleaf*
Fleabane, rough*	Smartweed, ladysthumb*
Florida pusley	Smartweed, Pennsylvania*
Foxtail*	Sowthistle, annual
Goatgrass, jointed*	Spanishneedles
Goosegrass	Speedwell, purslane*
Grain sorghum (milo)*	Sprangletop*
Groundsel, common*	Spurge, annual
Hemp sesbania	Spurge, prostrate*
Henbit	Spurge, spotted*
Horseweed/Marestail	Spurry, umbrella*
( <i>Conyza canadensis</i> )	Starthistle, yellow
Itchgrass*	Stinkgrass*
Johnsongrass, seedling	Sunflower*
Junglerice	Teaweed/Prickly sida
Knotweed	Texas panicum*
Kochia	Velvetleaf
Lamb's-quarters*	Virginia copperleaf
Little barley*	Virginia pepperweed*
London rocket*	Wheat*
Mayweed	Wild oats*
Medusahead*	Witchgrass*
Morningglory ( <i>Ipomoea spp</i> )	Woolly cupgrass*
Mustard, blue*	Yellow rocket

\*When using field broadcast equipment (aerial applications or boom sprayers using flat-fan nozzles) these species will be controlled or partially controlled using 1 pint of this product per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Use nozzles that ensure thorough coverage of foliage and treat when weeds are in an early growth stage.

## 9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the recommended range.

Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment. When using hand-held equipment for low-volume directed spot treatments, apply a 5- to 10-percent solution of this product.

Allow 7 or more days after application before tillage.

Weed Species	Rate (QT/A)	Hand-Held % Solution
Alfalfa*	1	2
Alligatorweed*	4	1.5
Anise (fennel)	2 - 4	1 - 2
Bahiagrass	3 - 5	2
Beachgrass, European ( <i>Ammophila arenaria</i> )	—	5
Bentgrass*	1.5	2
Bermudagrass	5	2
Bermudagrass, water (knotgrass)	1.5	2
Bindweed, field	4 - 5	2
Bluegrass, Kentucky	2	2
Blueweed, Texas	4 - 5	2
Brackenfern	3 - 4	1 - 1.5
Bromegrass, smooth	2	2
Bursage, woolly-leaf	—	2

Weed Species	Rate (QT/A)	Hand-Held % Solution
Canarygrass, reed	2 - 3	2
Cattail	3 - 5	2
Clover, red, white	3 - 5	2
Cogongrass	3 - 5	2
Dallisgrass	3 - 5	2
Dandelion	3 - 5	2
Dock, curly	3 - 5	2
Dogbane, hemp	4	2
Fescue (except tall)	3 - 5	2
Fescue, tall	1 - 3	2
German ivy	2 - 4	1 - 2
Guineagrass	3	1
Horsenettle	3 - 5	2
Horseradish	4	2
Iceplant	2	1.5 - 2
Jerusalem artichoke	3 - 5	2
Johnsongrass	2 - 3	1
Kikuyugrass	2 - 3	2
Knapweed	4	2
Lantana	—	1 - 1.25
Lespedeza	3 - 5	2
Milkweed, common	3	2
Muhly, wirestem	2	2
Mullein, common	3 - 5	2
Napiergrass	3 - 5	2
Nightshade, silverleaf	2	2
Nutsedge, purple, yellow	3	1 - 2
Orchardgrass	2	2
Pampasgrass	3 - 5	1.5 - 2
Paragrass	3 - 5	2
Pepperweed, perennial	4	2
Phragmites*	3 - 5	1 - 2
Poison hemlock	2 - 4	1 - 2
Quackgrass	2 - 3	2
Redvine*	2	2
Reed, giant	4 - 5	2
Ryegrass, perennial	2 - 3	1
Smartweed, swamp	3 - 5	2
Spurge, leafy*	—	2
Sweet potato, wild*	—	2
Thistle, artichoke	2 - 3	1 - 2
Thistle, Canada	2 - 3	2
Timothy	2 - 3	2
Torpedograss*	4 - 5	2
Trumpet creeper*	2 - 3	2
Vaseygrass	3 - 5	2
Velvetgrass	3 - 5	2
Wheatgrass, western	2 - 3	2

\*Partial control

## 9.3 Woody Brush and Trees

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment. When using hand-held equipment for low volume directed-spray spot treatments, apply a 5- to 10-percent solution of this product.

Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

Weed Species	Broadcast Rate (QT/A)	Hand-Held Spray-to-Wet % Solution
Alder	3 - 4	1 - 1.5
Ash*	2 - 5	1 - 2
Aspen, quaking	2 - 3	1 - 1.5
Bearclover (Bearmat)*	2 - 5	1 - 2
Beech*	2 - 5	1 - 2
Birch	2	1
Blackberry	3 - 4	1 - 1.5
Blackgum	2 - 5	1 - 2
Bracken	2 - 5	1 - 2
Broom; French, Scotch	2 - 5	1.5 - 2
Buckwheat, California*	2 - 4	1 - 2
Cascara*	2 - 5	1 - 2
Catsclaw*	—	1 - 1.5
Ceanothus*	2 - 5	1 - 2
Chamise*	2 - 5	1
Cherry; bitter, black, pin	2 - 3	1 - 1.5
Coyote brush	3 - 4	1.5 - 2
Deerweed	2 - 5	1
Dogwood*	2 - 5	1 - 2
Elderberry	2	1
Elm*	2 - 5	1 - 2
Eucalyptus	—	2
Gorse*	2 - 5	1 - 2
Hasardia*	2 - 4	1 - 2
Hawthorn	2 - 3	1 - 1.5
Hazel	2	1
Hickory*	2 - 5	1 - 2
Honeysuckle	3 - 4	1 - 1.5
Hornbeam, American*	2 - 5	1 - 2
Kudzu	4	2
Locust, black*	2 - 4	1 - 2
Madrone resprouts*	—	2
Manzanita*	2 - 5	1 - 2
Maple, red	2 - 4	1 - 1.5
Maple, sugar	—	1 - 1.5
Monkey flower*	2 - 4	1 - 2
Oak; black, white*	2 - 4	1 - 2
Oak, post	3 - 4	1 - 1.5
Oak, northern pin	2 - 4	1 - 1.5
Oak, scrub*	2 - 4	1 - 1.5
Oak, southern red	2 - 3	1 - 1.5
Peppertree, Brazilian (Florida holly)*	2 - 5	1 - 2
Persimmon*	2 - 5	1 - 2
Pine	2 - 5	1 - 2
Poison ivy	4 - 5	2
Poison oak	4 - 5	2
Poplar, yellow*	2 - 5	1 - 2
Redbud, eastern	2 - 5	1 - 2
Rose, multiflora	2	1
Russian olive*	2 - 5	1 - 2
Sage, black	2 - 4	1
Sage, white*	2 - 4	1 - 2
Sage brush, California	2 - 4	1
Salmonberry	2	1
Saltcedar*	2 - 5	1 - 2
Sassafras*	2 - 5	1 - 2
Sourwood*	2 - 5	1 - 2
Sumac; laurel, poison, smooth, sugarbush, winged*	2 - 4	1 - 2
Sweetgum	2 - 3	1 - 1.5
Swordfern*	2 - 5	1 - 2
Tallowtree, Chinese	—	1
Tan oak resprouts*	—	2
Thimbleberry	2	1
Tobacco, tree*	2 - 4	1 - 2
Toyon*	—	2
Trumpet creeper	2 - 3	1 - 1.5

Weed Species	Broadcast Rate (QT/A)	Hand-Held Spray-to-Wet % Solution
Vine maple*	2 - 5	1 - 2
Virginia creeper	2 - 5	1 - 2
Waxmyrtle, southern*	2 - 5	1 - 2
Willow	3	1
Yerbasanta, California*	—	2

\*Partial control

## 10.0 LIMIT OF WARRANTY AND LIABILITY

Monsanto Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Outrider, Ranger PRO, and Monsanto & Vine symbol are trademarks of Monsanto Technology LLC.

All other trademarks are the property of their respective owner.

This product is protected by U.S. Patent Nos. 5,683,958; 5,703,015; 6,063,733; 6,121,199; 6,121,200. No license granted under any non-U.S. patent(s).

EPA Reg. No. 524-517

In case of an emergency involving this product, or for medical assistance,  
Call Collect, day or night, (314) 694-4000.

Packed For:  
MONSANTO COMPANY  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI, 63167 USA  
©2007



# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

1/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

### SECTION 1: IDENTIFICATION OF THE SUBSTANCE/MIXTURE AND OF THE COMPANY/UNDERTAKING

#### Product identifier

Trade name	RANGER PRO® HERBICIDE
Product code (UVP)	86775093
SDS Number	102000037601
EPA Registration No.	524-517

#### Relevant identified uses of the substance or mixture and uses advised against

Identified uses	Herbicide
Restrictions on use	See product label for restrictions.
Information on supplier	
Supplier	Bayer Environmental Science A division of Bayer CropScience LP 500 Centregreen Way, Suite 400 Cary, NC 27513 USA
Responsible Department	Email: SDSINFO.BCS-NA@bayer.com
Emergency telephone no.	
Emergency Telephone Number (24hr/ 7 days)	1-800-334-7577
Product Information Telephone Number	1-800-331-2867

### SECTION 2: HAZARDS IDENTIFICATION

Classification in accordance with regulation HCS 29CFR §1910.1200  
Acute toxicity(Inhalation): Category 4

Labelling in accordance with regulation HCS 29CFR §1910.1200



Signal word: Warning

Hazard statements  
Harmful if inhaled.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

2/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

### Precautionary statements

Avoid breathing mist/ vapours/ spray.  
Use only outdoors or in a well-ventilated area.  
IF INHALED: Remove person to fresh air and keep comfortable for breathing.  
Call a POISON CENTER/doctor/physician if you feel unwell.

### Hazards Not Otherwise Classified (HNOC)

No physical hazards not otherwise classified.  
No health hazards not otherwise classified.

## SECTION 3: COMPOSITION/INFORMATION ON INGREDIENTS

Hazardous Component Name	CAS-No.	Concentration % by weight
Isopropylamine salt of glyphosate	38641-94-0	41.0
Surfactant blend (proprietary)		>=5.0 – <=10.0

The specific chemical identity and/or concentration range is being withheld because it is trade secret information.

## SECTION 4: FIRST AID MEASURES

### Description of first aid measures

<b>General advice</b>	When possible, have the product container or label with you when calling a poison control center or doctor or going for treatment.
<b>Inhalation</b>	Move to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a physician or poison control center immediately.
<b>Skin contact</b>	Wash off immediately with plenty of water for at least 15 minutes. Take off contaminated clothing and shoes immediately. Call a physician or poison control center immediately.
<b>Eye contact</b>	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a physician or poison control center immediately.
<b>Ingestion</b>	Call a physician or poison control center immediately. Rinse out mouth and give water in small sips to drink. DO NOT induce vomiting unless directed to do so by a physician or poison control center. Never give anything by mouth to an unconscious person. Do not leave victim unattended.

### Most important symptoms and effects, both acute and delayed

<b>Symptoms</b>	To date no symptoms are known.
<b>Indication of any immediate medical attention and special treatment needed</b>	
<b>Risks</b>	This product is not a cholinesterase inhibitor.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

3/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

**Treatment** Treatment with atropine and oximes is not indicated. Appropriate supportive and symptomatic treatment as indicated by the patient's condition is recommended.

---

### SECTION 5: FIREFIGHTING MEASURES

#### Extinguishing media

**Suitable** Use water spray, alcohol-resistant foam, dry chemical or carbon dioxide.

**Unsuitable** High volume water jet

**Special hazards arising from the substance or mixture** In the event of fire the following may be released: Carbon monoxide (CO), Carbon dioxide (CO<sub>2</sub>), Nitrogen oxides (NO<sub>x</sub>), Oxides of phosphorus

#### Advice for firefighters

**Special protective equipment for firefighters** In the event of fire and/or explosion do not breathe fumes. Firefighters should wear NIOSH approved self-contained breathing apparatus and full protective clothing. Equipment should be thoroughly decontaminated after use.

**Further information** Keep out of smoke. Fight fire from upwind position. Cool closed containers exposed to fire with water spray. Do not allow run-off from fire fighting to enter drains or water courses.

**Flash point** does not flash

**Auto-ignition temperature** No data available

**Lower explosion limit** Not applicable

**Upper explosion limit** Not applicable

**Explosivity** Not explosive

---

### SECTION 6: ACCIDENTAL RELEASE MEASURES

#### Personal precautions, protective equipment and emergency procedures

**Precautions** Use personal protective equipment. Keep unauthorized people away. Avoid contact with spilled product or contaminated surfaces.

#### Methods and materials for containment and cleaning up

**Methods for cleaning up** Soak up with inert absorbent material (e.g. sand, silica gel, acid binder, universal binder, sawdust). Collect and transfer the product into a properly labelled and tightly closed container. Keep in suitable, closed containers for disposal. Clean contaminated floors and objects thoroughly, observing environmental regulations.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

4/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

---

<b>Additional advice</b>	Use personal protective equipment. If the product is accidentally spilled, do not allow to enter soil, waterways or waste water canal. Do not allow product to contact non-target plants.
<b>Reference to other sections</b>	Information regarding safe handling, see section 7. Information regarding personal protective equipment, see section 8. Information regarding waste disposal, see section 13.

---

### SECTION 7: HANDLING AND STORAGE

#### Precautions for safe handling

**Advice on safe handling** Avoid contact with skin, eyes and clothing. Ensure adequate ventilation.

**Hygiene measures** Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, using the toilet or applying cosmetics.  
Remove Personal Protective Equipment (PPE) immediately after handling this product. Remove soiled clothing immediately and clean thoroughly before using again. Wash thoroughly and put on clean clothing. Keep working clothes separately. Garments that cannot be cleaned must be destroyed (burnt).

#### Conditions for safe storage, including any incompatibilities

**Requirements for storage areas and containers** Store in original container. Store in a cool, dry place and in such a manner as to prevent cross contamination with other crop protection products, fertilizers, food, and feed. Store in a place accessible by authorized persons only. Reacts with galvanised steel or unlined mild steel to produce hydrogen, a highly flammable gas that could explode. Protect from freezing. Partial crystallization may occur on prolonged storage below the minimum storage temperature. Freezing will affect the physical condition but will not damage the material. Thaw and mix before using.

**Advice on common storage** Keep away from food, drink and animal feedingstuffs.

---

### SECTION 8: EXPOSURE CONTROLS/PERSONAL PROTECTION

#### Control parameters

No known occupational limit values.

#### Exposure controls

##### Personal protective equipment

In normal use and handling conditions please refer to the label and/or leaflet. In all other cases the following recommendations would apply.

##### Respiratory protection

When respirators are required, select NIOSH approved equipment based on actual or potential airborne concentrations and in accordance with the appropriate regulatory standards and/or industry recommendations.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

5/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

<b>Hand protection</b>	Please observe the instructions regarding permeability and breakthrough time which are provided by the supplier of the gloves. Also take into consideration the specific local conditions under which the product is used, such as the danger of cuts, abrasion, and the contact time. Chemical-resistant gloves (barrier laminate, butyl rubber, nitrile rubber or Viton) Wash gloves when contaminated. Dispose of when contaminated inside, when perforated or when contamination on the outside cannot be removed. Wash hands frequently and always before eating, drinking, smoking or using the toilet.
<b>Eye protection</b>	Use tightly sealed goggles and face protection.
<b>Skin and body protection</b>	Wear long-sleeved shirt and long pants and shoes plus socks.
<b>General protective measures</b>	Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and warm/tepid water. Keep and wash PPE separately from other laundry.

## SECTION 9. PHYSICAL AND CHEMICAL PROPERTIES

### Information on basic physical and chemical properties

<b>Form</b>	Liquid, clear
<b>Colour</b>	light yellow to amber
<b>Odour</b>	sweet
<b>Odour Threshold</b>	No data available
<b>pH</b>	4.4 - 5.0 (8 %) (23 °C) (deionized water)
<b>Melting point/range</b>	No data available
<b>Boiling Point</b>	No data available
<b>Flash point</b>	does not flash
<b>Flammability</b>	No data available
<b>Auto-ignition temperature</b>	No data available
<b>Minimum ignition energy</b>	Not applicable
<b>Self-accelarating decomposition temperature (SADT)</b>	No data available
<b>Upper explosion limit</b>	Not applicable
<b>Lower explosion limit</b>	Not applicable
<b>Vapour pressure</b>	No significant volatility.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

6/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

<b>Evaporation rate</b>	No data available
<b>Relative vapour density</b>	No data available
<b>Relative density</b>	1.162 (20 °C)
<b>Density</b>	1.17 g/cm <sup>3</sup> (20 °C)
<b>Water solubility</b>	completely miscible
<b>Partition coefficient: n-octanol/water</b>	Glyphosate: log Pow: -3.2
<b>Viscosity, dynamic</b>	No data available
<b>Viscosity, kinematic</b>	No data available
<b>Oxidizing properties</b>	No data available
<b>Explosivity</b>	Not explosive
<b>Other information</b>	Further safety related physical-chemical data are not known.

### SECTION 10: STABILITY AND REACTIVITY

#### Reactivity

<b>Thermal decomposition</b>	Stable under normal conditions.
<b>Chemical stability</b>	Stable under recommended storage conditions.
<b>Possibility of hazardous reactions</b>	Reacts with galvanised steel or unlined mild steel to produce hydrogen, a highly flammable gas that could explode.
<b>Conditions to avoid</b>	Extremes of temperature and direct sunlight.
<b>Incompatible materials</b>	Galvanised steel, Unlined mild steel
<b>Hazardous decomposition products</b>	No decomposition products expected under normal conditions of use.

### SECTION 11: TOXICOLOGICAL INFORMATION

<b>Exposure routes</b>	Skin contact, Eye contact, Inhalation
<b>Immediate Effects</b>	
<b>Eye</b>	May cause eye irritation.
<b>Skin</b>	Not expected to produce significant adverse effects when recommended use instructions are followed.
<b>Ingestion</b>	Not expected to produce significant adverse effects when recommended use instructions are followed.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

7/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

<b>Inhalation</b>	May be harmful if inhaled.
<b>Information on toxicological effects</b>	
<b>Acute oral toxicity</b>	LD50 (Rat) 5,108 mg/kg Test conducted with a similar formulation.
<b>Acute inhalation toxicity</b>	LC50 (Rat) 2.9 mg/l Exposure time: 4 h Determined in the form of liquid aerosol. Test conducted with a similar formulation.
<b>Acute dermal toxicity</b>	LD50 (Rat) > 5,000 mg/kg Test conducted with a similar formulation. No deaths
<b>Skin corrosion/irritation</b>	No skin irritation (Rabbit) Test conducted with a similar formulation.
<b>Serious eye damage/eye irritation</b>	Slight irritant effect - does not require labelling. (Rabbit) Test conducted with a similar formulation.
<b>Respiratory or skin sensitisation</b>	Skin: Non-sensitizing. (Guinea pig) OECD Test Guideline 406, Buehler test Test conducted with a similar formulation.

### Assessment STOT Specific target organ toxicity – single exposure

Glyphosate: Based on available data, the classification criteria are not met.

### Assessment STOT Specific target organ toxicity – repeated exposure

Glyphosate did not cause specific target organ toxicity in experimental animal studies.

### Assessment mutagenicity

Glyphosate was not mutagenic or genotoxic in a battery of in vitro and in vivo tests.

### Assessment carcinogenicity

Glyphosate was not carcinogenic in lifetime feeding studies in rats and mice.  
Important comment to IARC Listing: Our expert opinion is that classification as a carcinogen is not warranted.

### ACGIH

None.

### NTP

None.

### IARC

Isopropylamine salt of glyphosate                      38641-94-0                      Overall evaluation: 2A

### OSHA

None.

### Assessment toxicity to reproduction

Glyphosate did not cause reproductive toxicity in a two-generation study in rats.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

8/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

### Assessment developmental toxicity

Glyphosate did not cause developmental toxicity in rats and rabbits.

### Aspiration hazard

Based on available data, the classification criteria are not met.

### Further information

The toxicological data refer to a similar formulation.

---

## SECTION 12: ECOLOGICAL INFORMATION

<b>Toxicity to fish</b>	LC50 (Oncorhynchus mykiss (rainbow trout)) 5.4 mg/l static test; Exposure time: 96 h Test conducted with a similar formulation.
<b>Chronic toxicity to fish</b>	Oncorhynchus mykiss (rainbow trout) flow-through test NOEC: $\geq 9.63$ mg/l The value mentioned relates to the active ingredient glyphosate.
<b>Toxicity to aquatic invertebrates</b>	EC50 (Daphnia magna (Water flea)) 11 mg/l static test; Exposure time: 48 h Test conducted with a similar formulation.
<b>Chronic toxicity to aquatic invertebrates</b>	EC50 (Daphnia magna (Water flea)): 12.5 mg/l Exposure time: 21 d The value mentioned relates to the active ingredient glyphosate.
<b>Toxicity to aquatic plants</b>	EbC50 (Raphidocelis subcapitata (freshwater green alga)) 12.4 mg/l static test; Exposure time: 72 h Test conducted with a similar formulation.  NOEC (Raphidocelis subcapitata (freshwater green alga)) 6.3 mg/l static test; Exposure time: 72 h Test conducted with a similar formulation.
<b>Biodegradability</b>	Glyphosate: Not rapidly biodegradable
<b>Koc</b>	Glyphosate: Koc: 6920
<b>Bioaccumulation</b>	Glyphosate: Does not bioaccumulate.
<b>Mobility in soil</b>	Glyphosate: Immobile in soil
<b>Results of PBT and vPvB assessment</b>	
<b>PBT and vPvB assessment</b>	Glyphosate: This substance is not considered to be persistent, bioaccumulative and toxic (PBT). This substance is not considered to be very persistent and very bioaccumulative (vPvB).
<b>Additional ecological</b>	No further ecological information is available.

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

9/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

### information

**Environmental precautions** Apply this product as specified on the label.  
Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.  
Do not contaminate surface or ground water by cleaning equipment or disposal of wastes, including equipment wash water.  
Retain and dispose of contaminated wash water.

---

## SECTION 13: DISPOSAL CONSIDERATIONS

### Waste treatment methods

**Product** It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow container label instructions and applicable local guidelines.  
Do not contaminate water, food, or feed by disposal.  
Follow all local/regional/national/international regulations.

**Contaminated packaging** Follow advice on product label and/or leaflet.  
Do not re-use empty containers.  
Triple rinse containers.  
Puncture container to avoid re-use.  
Completely empty container into application equipment, then dispose of empty container in a sanitary landfill, by incineration or by other procedures approved by state/provincial and local authorities.  
If burned, stay out of smoke.

**RCRA Information** Characterization and proper disposal of this material as a special or hazardous waste is dependent upon Federal, State and local laws and are the user's responsibility. RCRA classification may apply.

---

## SECTION 14: TRANSPORT INFORMATION

According to national and international transport regulations this material is not classified as dangerous goods / hazardous material.

Freight Classification: COMPOUNDS, TREE OR WEED KILLING, N.O.I. other than poison, HAVING A DENSITY OF 20 LBS OR GREATER PER CUBIC FOOT

---

## SECTION 15: REGULATORY INFORMATION

EPA Registration No. 524-517

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

10/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

### US Federal Regulations

#### TSCA list

Water 7732-18-5  
Polyethylene glycol 25322-68-3

#### US. Toxic Substances Control Act (TSCA) Section 12(b) Export Notification (40 CFR 707, Subpt D)

No export notification needs to be made.

#### SARA Title III - Section 302 - Notification and Information

Not applicable.

#### SARA Title III - Section 313 - Toxic Chemical Release Reporting

None.

### US States Regulatory Reporting

#### CA Prop65

This product does not contain any substances known to the State of California to cause cancer.

This product does not contain any substances known to the State of California to cause reproductive harm.

### US State Right-To-Know Ingredients

Polyethylene glycol 25322-68-3 MN

### Environmental

#### CERCLA

None.

#### Clean Water Section 307(a)(1)

None.

#### Safe Drinking Water Act Maximum Contaminant Levels

None.

### EPA/FIFRA Information:

This chemical is a pesticide product registered by the Environmental Protection Agency and is subject to certain labeling requirements under federal pesticide law. These requirements differ from the classification criteria and hazard information required for safety data sheets, and for workplace labels of non-pesticide chemicals. Following is the hazard information required on the pesticide label:

**Signal word:** Caution!

**Hazard statements:** Causes eye irritation.

## SECTION 16: OTHER INFORMATION

### Abbreviations and acronyms

49CFR	Code of Federal Regulations, Title 49
ACGIH	US. ACGIH Threshold Limit Values
ATE	Acute toxicity estimate
CAS-Nr.	Chemical Abstracts Service number
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act

# SAFETY DATA SHEET



## RANGER PRO® HERBICIDE

Version 1.0 / USA  
102000037601

11/11  
Revision Date: 08/06/2020  
Print Date: 08/17/2020

EINECS	European inventory of existing commercial substances
ELINCS	European list of notified chemical substances
IARC	International Agency for Research on Cancer
IATA	International Air Transport Association
IMDG	International Maritime Dangerous Goods
N.O.S.	Not otherwise specified
NTP	US. National Toxicology Program (NTP) Report on Carcinogens
OECD	Organization for Economic Co-operation and Development
TDG	Transportation of Dangerous Goods
TWA	Time weighted average
UN	United Nations
WHO	World health organisation

### NFPA 704 (National Fire Protection Association):

Health - 1      Flammability - 1      Instability - 1      Others - none

### HMIS (Hazardous Materials Identification System, based on the Third Edition Ratings Guide)

Health - 2      Flammability - 1      Physical Hazard - 1      PPE -

0 = minimal hazard, 1 = slight hazard, 2 = moderate hazard, 3 = severe hazard, 4 = extreme hazard

**Reason for Revision:** New Safety Data Sheet.

**Revision Date:** 08/06/2020

This information is provided in good faith but without express or implied warranty. The customer assumes all responsibility for safety and use not in accordance with label instructions. The product names are registered trademarks of Bayer.

# SPECIMEN

Pendimethalin

Group

3

Herbicide

# Pendulum® AquaCap

## Herbicide

**For use as a preemergence weed control herbicide in turfgrass, landscape or grounds maintenance, noncropland areas, and ornamental production**

**Active Ingredient:**

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine . . . . . 38.7%

**Other Ingredients:** . . . . . 61.3%

**Total:** . . . . . 100.0%

1 gallon contains 3.8 lbs of microencapsulated pendimethalin in an aqueous carrier.

EPA Reg. No. 241-416

EPA Est. No.

### KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

**In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

**Net Contents:**

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOTLINE</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of an emergency endangering life or property involving this product, call day or night, 1-800-832-HELP (4357).	

## **Precautionary Statements**

### **Hazards to Humans and Domestic Animals**

**CAUTION.** Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

### **Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

### **Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.240)(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## **USER SAFETY RECOMMENDATIONS**

### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## **Environmental Hazards**

This product is toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

**Non-target Organism Advisory Statement:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## **Directions For Use**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at time of herbicide application.

**DO NOT** apply this product through any type of irrigation system.

BASF Corporation does not recommend or authorize the use of this product in manufacturing, processing or preparing custom blends with other products for application to turf or ornamentals.

**DO NOT** apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application.

For requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

**DO NOT** apply **Pendulum® AquaCap herbicide** in greenhouses, shadehouses, or other enclosed structures.

Not for use for commercial seed production.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

## NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**DO NOT** enter treated areas without protective clothing until sprays have dried.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL OR CROP INJURY.**

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

### Pesticide Storage

**DO NOT** store below 15° F. Extended storage at temperatures below 15° F can result in the formation of crystals on the bottom of container. If crystallization does occur, store the container on its side at room temperature (70° F) and rock occasionally until crystals dissolve.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### Container Handling

**Nonrefillable Container. DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Triple rinse containers too large to shake (capacity > 5 gallons) as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

*(continued)*

## STORAGE AND DISPOSAL *(continued)*

### Container Handling *(continued)*

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

**Triple rinse as follows:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

## Product Information

**Pendulum® AquaCap herbicide** controls most annual grasses and certain broadleaf weeds as they germinate. When susceptible weeds germinate in the treated area, they contact the herbicide and both shoot and root growth stops. Translocation of the herbicide within the plant is limited. Affected weeds die shortly after growth is stopped, usually before emergence from the soil.

### Mode of Action

Pendimethalin, the active ingredient in **Pendulum AquaCap**, is a **Group 3 (WSSA)/Group K<sub>1</sub> (HRAC)** herbicide belonging to the dinitroaniline chemistry class. **Pendulum AquaCap** is a meristematic inhibitor that interferes with meristematic plant cell division or mitosis inhibiting germinating seedling growth.

## Herbicide Resistance Management

While weed resistance to **Group 3** herbicides is infrequent, populations of resistant biotypes are known to exist. Weeds resistant to **Group 3** herbicides should be managed using herbicide(s) from a different group (mode or site of action) that are effective against the target weeds. Resistance management should be part of a diversified weed control strategy that integrates chemical, cultural, and mechanical (tillage) control tactics. Consult your local BASF representative, state cooperative extension service, professional consultants, or other qualified authority to determine appropriate actions if you suspect resistant weeds.

### Chemical Control

- Start clean with mechanical control measure or an effective burndown herbicide program.
- **DO NOT** rely on a single herbicide site of action for weed control.
- Follow labeled application rate and weed growth stage specifications.
- Avoid application of herbicides with the same site of action more than twice a season.
- Use tank mixes and sequential applications with other herbicides possessing different sites of action that are also effective on the target weeds.

### Scouting and Containment

- Scout treated areas after herbicide application to identify areas where weed control was ineffective.
- Control weed escapes with herbicides possessing a different site of action or use a mechanical control measure. Weed escapes should not be allowed to reproduce by seed or to proliferate vegetatively.
- Contact your **Pendulum AquaCap** supplier and/or your local BASF representative to report weed escapes.
- Clean equipment before moving to a different treatment area to avoid spread of resistant weeds.

### Weeds Controlled

**Pendulum AquaCap** will not control established weeds. If weeds germinate before herbicide activation, shallow cultivate to destroy existing weeds or, where practical, remove by hand. When cultivating for any reason, it should be shallow. Use **Pendulum AquaCap** with herbicides registered for postemergence application (i.e. **Roundup® herbicide** or **Finale® herbicide**) for the control of established weeds. **DO NOT** apply sprays containing **Roundup** or **Finale** over the top of desirable plants. A **Pendulum AquaCap** treatment may be followed by any registered herbicide to control weeds not listed on the **Pendulum AquaCap** label.

The efficacy of **Pendulum AquaCap** will be improved if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. Erratic weed control may result if **Pendulum AquaCap** is not activated by rainfall or irrigation within 30 days.

The following grass and broadleaf weeds are controlled by preemergence treatments of **Pendulum AquaCap** at the specified rates.

**Table 1. Weeds Controlled**

Common Name	Scientific Name
<b>Grasses</b>	
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bluegrass, annual	<i>Poa annua</i>
Crabgrass	<i>Digitaria</i> spp.
Crowfootgrass	<i>Dactyloctenium aegyptium</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria glauca</i>
Goosegrass	<i>Eleusine indica</i>
Itchgrass	<i>Rottboellia exaltata</i>
Johnsongrass (from seed)	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colona</i>
Lovegrass (from seed)	<i>Eragrostis</i> spp.
Panicum, browntop	<i>Panicum fasciculatum</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Sandbur, field	<i>Cenchrus incertus</i>
Signalgrass	<i>Brachiaria platyphylla</i>
Sprangletop, Mexican	<i>Leptochloa uninervia</i>
Sprangletop, red	<i>Leptochloa filiformis</i>
Witchgrass	<i>Panicum capillare</i>
Woolly cupgrass	<i>Eriochloa villosa</i>
<b>Broadleaf Weeds</b>	
Burweed, lawn	<i>Soliva pterosperma</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Clover, hop	<i>Trifolium procumbens</i>
Cudweed	<i>Gnaphalium</i> spp.
Evening primrose	<i>Oenothera biennis</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Filaree	<i>Erodium</i> spp.
Henbit	<i>Lamium amplexicaule</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters	<i>Chenopodium album</i>
Pigweed	<i>Amaranthus</i> spp.
Puncturevine	<i>Tribulus terrestris</i>
Purslane	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>

(continued)

**Table 1. Weeds Controlled** (continued)

Common Name	Scientific Name
<b>Broadleaf Weeds</b> (continued)	
Rocket, London	<i>Sisymbrium irio</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Speedwell, corn	<i>Veronica arvensis</i>
Spurge, annual	<i>Euphorbia</i> spp.
Spurge, prostrate	<i>Chamaesyce humistrata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>
Velvetleaf (Buttonweed)	<i>Abutilon theophrasti</i>

### Application Use Sites

Use **Pendulum® AquaCap herbicide** for preemergence control of grass and certain broadleaf weed species as they germinate **in any turfgrass site** (golf courses, lawns, sod farms and other turf areas) and **landscape ornamental maintenance areas**. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multifamily dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses, prairie grass areas, and sod farms.

**Pendulum AquaCap can be applied for general grounds maintenance** in areas such as parking lots, driveways and roadsides, alleyways, bike and jogging paths, vacant lots, buildings, stone gardens and gravel yards, markers and fence lines, and mulch beds. It may be used under asphalt or concrete treatments as part of a site preparation program.

Use **Pendulum AquaCap** for preemergence control of most annual grasses and certain broadleaf weeds as they germinate **in any noncropland area** such as railroad, utility, highway, and pipeline rights-of-way; highway guardrails, delineators, and sign posts; bridge abutments and approaches; utility substations; petroleum tank farms; pumping installations; storage areas; fence rows; windbreaks and shelterbelts; paved or gravel surfaces; and established wildflower plantings where weed control is desired.

**Pendulum AquaCap** can also be used **in bulb plantings, nonbearing fruit and nut tree nurseries, conifer and hardwood seedling nurseries, and tree plantations for site preparation and maintenance**.

Applications can be made, but are not limited to, plant species listed on this label such as trees, shrubs, groundcovers, perennials, bulbs, ornamental grasses, and bedding plants.

**Pendulum AquaCap** can be used **in and around field, liner, and container ornamental production**.

---

## Application Instructions

---

**Pendulum® AquaCap herbicide** will not control established weeds. Therefore, areas to be treated should be free of established weeds at the time of treatment, or **Pendulum AquaCap** may be used with herbicides registered for postemergence use in managed turf sites, landscape ornamentals, and in other noncropland areas. Consult the labels of those herbicides for suggested treatments, rates, and precautions or restrictions for use in these areas. The efficacy of **Pendulum AquaCap** will improve if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. If **Pendulum AquaCap** is not activated by rainfall or irrigation within 30 days, erratic weed control may result.

Applied according to label directions and under normal growing conditions, **Pendulum AquaCap** or **Pendulum AquaCap** tank mix combinations will not cause crop injury. Overapplication can result in crop stand loss, crop injury, or soil residues. Uneven application can decrease weed control or cause crop injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken seedlings and plants and increase the possibility of plant damage from **Pendulum AquaCap**.

---

## Mixing Instructions

---

**Pendulum AquaCap** may be applied in a tank mix or a sequential application with other herbicides registered for use in a given crop. Refer to the companion label for weeds controlled in addition to **Pendulum AquaCap** alone.

When using tank mixtures or sequential applications with **Pendulum AquaCap**, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

Fill tank 1/2 to 3/4 full with clean water or liquid fertilizer and agitate. Before mixing **Pendulum AquaCap** or **Pendulum AquaCap** tank mixtures in liquid fertilizer, refer to appropriate label sections for recommended uses in liquid fertilizer, application instructions, and compatibility determinations.

### Pendulum AquaCap Alone

When using **Pendulum AquaCap** alone, add **Pendulum AquaCap** to the partially filled tank while agitating; then fill the remainder of the tank with water or liquid fertilizer.

### Pendulum AquaCap Tank Mixes

Add the tank mixture ingredients in the following order:

1. **Wettable Powder (WP) formulations** - Make a slurry of the WP in water (1:2 ratio). Add the slurry slowly into the partially filled tank while agitating.

2. **Dry Flowable/Water Dispersible Granule (DF/WDG) formulations** - Add the granules to the partially filled tank while agitating. Make a slurry of the granules in water before adding to liquid fertilizer.
3. **Flowable (F) formulations** - Add the F formulation to the partially filled tank while agitating.
4. Add **Pendulum AquaCap** to the partially filled tank while agitating.
5. **Water-soluble Concentrate (WSC) formulations** - Add the WSC formulation to the partially filled tank while agitating.
6. **Emulsifiable Concentrate (EC) formulations** - Add the EC formulation to the partially filled tank while agitating.

Fill the remainder of the tank with water or liquid fertilizer while agitating.

Maintain continuous agitation while adding herbicides and until spraying is completed. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed.

## Backpack Sprayer

Begin with a clean spray tank. Fill the spray tank 1/2 full with clean water and add the required amount of **Pendulum AquaCap** to the sprayer. Cap sprayer and agitate to ensure mixing. Uncap sprayer and finish filling tank to desired level. Cap sprayer and agitate once again. During application it is desirable to agitate the mixture on occasion to ensure mixing. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential before spraying is resumed.

## Liquid Fertilizers

Before mixing, always test small quantities with a simple jar test. Add the required amount of **Pendulum AquaCap** to a half-filled spray tank while agitating; then add the fertilizer product. Complete filling spray tank to desired level.

---

## Spraying Instructions

---

Avoid unintentional contact of spray solution with driveways, stone, wood, or other porous surfaces. Rinse immediately with water to avoid staining. Avoid mechanically scrubbing until surface area is thoroughly rinsed. Treated turfgrass should be dry before entering to avoid staining onto nontreated surfaces.

## Ground Application (spray boom)

Uniformly apply with properly calibrated ground equipment in sufficient water per acre to uniformly treat the area with a spray pressure of 25 to 50 psi. Suggested spray volumes are 20 to 200 gpa for professional turfgrass, landscape and ornamental applications, and 10 to 200 gpa for all other noncrop applications such as roadsides, utility rights-of-way, or soft-residual bareground applications. Maintain continuous agitation during spraying with good mechanical or bypass agitation. Avoid overlaps that will increase rates above those specified.

## Ground Application (hand-held equipment)

Use **Table 2** or **Table 3** to determine the amount of **Pendulum® AquaCap herbicide** to apply per 1,000 square feet of treated area. The amount of water used for the application is not critical but should be sufficient for thorough coverage without runoff. Calibration of backpack or other hand-held equipment will vary with each operator. Determine the amount of water needed to treat 1,000 square feet before mixing the spray solution. Follow information in **Mixing Instructions** section of this label.

## Aerial Application

Uniformly apply in 5 or more gallons of water per acre.

### Spray Drift Management

#### Ground Application (spray boom)

- Applicators must only apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or plant canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

#### Ground Application (hand-held)

- Take precautions to minimize spray drift.

### Spray Drift Management

#### Aerial Application

- **DO NOT** release spray at a height greater than 10 feet above the ground or plant canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must not exceed 65% of the wingspan for fixed wing aircraft or 75% of the rotor diameter for helicopters. Otherwise, the boom length must not exceed 75% of the wingspan for fixed wing aircraft or 90% of the rotor diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply during temperature inversions.

## Spray Drift Advisories

The applicator is responsible for avoiding off-site spray drift. Be aware of nearby non-target sites and environmental conditions.

### Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray

drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### Boom Height - Ground Boom

For ground equipment, the boom should remain level with the plants and have minimal bounce.

### Release Height - Aircraft

Higher release heights increase the potential for spray drift.

### Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### Temperature Inversion

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### Wind

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

---

## Turfgrass

---

Use **Pendulum® AquaCap herbicide** for preemergence control of grasses and certain broadleaf weed species as they germinate in any turfgrass site (golf courses, lawns, sod farms and other turf areas) and landscape ornamental maintenance areas. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multifamily dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses, prairie grass areas, and sod farms.

The efficacy of **Pendulum AquaCap** will be improved if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. If **Pendulum AquaCap** is not activated by rainfall or irrigation within 30 days, erratic weed control may result.

To prevent establishment of weeds along the edges of treated area, it may be necessary to overlap the spray 3 to 6 inches onto sidewalks or driveways, etc., to ensure effective application rates in these especially vulnerable sites. Where temporary discoloration of pavement is to be avoided, **DO NOT** rub or scrub surface. Rinse area immediately using a heavy spray of water to avoid staining. Treated turfgrass should be dry before entering to avoid staining onto nontreated surfaces.

### Turfgrass Tank Mixes

**Pendulum AquaCap** can be mixed with postemergence herbicides to control emerged weeds in nonresidential turfgrass. For annual grass control, applications can be made with **Drive® 75 DF herbicide**, **Drive® XLR8 herbicide**, or MSMA to control emerged weeds.

Broadleaf weeds can be controlled using **Trimec® herbicide**, **Three-Way™ herbicide**, 2-4,D and other similar products.

Before tank mixing, use a simple jar test to ensure compatibility of herbicides.

Refer to manufacturer's labels for specific use directions, precautions, and limitations before tank mixing with **Pendulum AquaCap**. Follow those that are most restrictive.

### Turfgrass Restrictions

- Use on well-established turfgrass with a dense and uniform stand. On turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery before application.
- On newly planted areas, application should not be made until the turfgrass has filled in and has been mowed at least four times. Applications made to overseeded warm-season turfgrass may cause thinning or injury of the overseeded species.
- **DO NOT** use on bentgrass or *Poa annua* greens and tees or injury may occur.

- Delay reseeding or winter overseeding treated turfgrass for at least three (3) months following the last **Pendulum AquaCap** application.
- Delay sprigging turfgrass for five (5) months after application.

**Table 2. Pendulum® AquaCap herbicide Residential, Golf Course, Commercial, and Other Nonresidential Turfgrass Uses for Preemergence Weed Control<sup>1</sup>**

Cool Season Turfgrass	Weed	Product per 1,000 sq ft (fl ozs)	Product per acre (pts)	Comment	
Bluegrass, Kentucky Fescue, fine Fescue, tall Ryegrass, perennial	barnyardgrass crabgrass evening primrose fall panicum foxtail hop clover knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	<b>All Turf Uses:</b>		Apply a repeat application of 2.2 to 3.1 pts/A (0.86 to 1.1 fl ozs/1,000 sq ft) after 5 to 8 weeks for extended control or where heavy weed infestations are expected.	
		1.1 to 1.6	3.1 to 4.2		
		Initial application before weed germination in spring			
	goosegrass	<b>Residential and Sod Farm Turf Uses Only<sup>2</sup>:</b>		Apply a repeat application of 3.1 pts/A (1.1 fl ozs/1,000 sq ft) if the lower rate was used initially or for extended goosegrass control after 5 to 8 weeks.	
		1.1 to 1.6	3.1 to 4.2		
		<b>Golf Course, Commercial and Other Nonresidential Turf Uses Only:</b>			
		1.1 to 2.3	3.1 to 6.3		
		Initial application before weed germination in spring			
	chickweed corn speedwell cudweed henbit lawn burweed <i>Poa annua</i>	<b>All Turf Uses:</b>		Apply in late summer or early fall before weed germination. Apply a repeat application of 3.1 to 4.2 pts/A (1.1 to 1.6 fl ozs/1,000 sq ft) after 5 to 8 weeks for extended <i>Poa annua</i> control.	
		1.1 to 1.6	3.1 to 4.2		
	Bentgrass or established <i>Poa annua</i> <sup>3</sup> (1/2-inch high or taller)	barnyardgrass crabgrass evening primrose fall panicum foxtail hop clover knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	<b>All Turf Uses (Non-greens and Tees):</b>		Apply a repeat application of 2.2 to 3.1 pts/A (0.86 to 1.1 fl ozs/1,000 sq ft) after 5 to 8 weeks for extended control or where heavy weed infestations are expected.
			1.1	3.1	
Initial application before weed germination in spring					
goosegrass		<b>All Turf Uses (Non-greens and Tees):</b>		Apply a repeat application of 3.1 pts/A (1.1 fl ozs/1,000 sq ft) for extended goosegrass control after 5 to 8 weeks.	
		1.1	3.1		
		Initial application before weed germination in spring			
chickweed corn speedwell cudweed henbit lawn burweed <i>Poa annua</i>		<b>All Turf Uses (Non-greens and Tees):</b>		Apply in late summer or early fall before weed germination.	
		1.1 to 1.6	3.1 to 4.2		

(continued)

**Table 2. Pendulum® AquaCap herbicide Residential, Golf Course, Commercial, and Other Nonresidential Turfgrass Uses for Preemergence Weed Control<sup>1</sup> (continued)**

Warm Season Turfgrass	Weed	Product per 1,000 sq ft (fl ozs)	Product per acre (pts)	Comment
Bahigrass Bermudagrass Buffalograss Centipedegrass Fescue, tall <i>Paspalum</i> , seashore St. Augustinegrass Zoysiagrass	barnyardgrass crabgrass evening primrose fall panicum foxtail hop clover knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	<b>Residential and Sod Farm Turf Uses Only:</b>		Apply a repeat application of 2.2 to 3.1 pts/A (0.86 to 1.1 fl ozs/1,000 sq ft) after 5 to 8 weeks if necessary.
		1.1 to 1.6	3.1 to 4.2	
		<b>Golf Course, Commercial and Other Nonresidential Turf Uses Only:</b>		
		1.1 to 2.3	3.1 to 6.3	
		Initial application before weed germination in spring		
	goosegrass	<b>All Turf Uses (Non-greens and Tees):</b>		An additional application of 3.1 pts/A (1.1 fl ozs/1,000 sq ft) may be made for extended goosegrass control 8 weeks after the second application.
		1.1	3.1	
		Apply before weed germination in spring. Make a second application at 3.1 pts/A (1.1 fl ozs/1,000 sq ft) 5 to 8 weeks later.		
	chickweed corn speedwell cudweed henbit lawn burweed <i>Poa annua</i>	<b>All Turf Uses:</b>		Apply in late summer or early fall before weed germination. Apply a repeat application of 3.1 to 4.2 pts/A (1.1 to 1.6 fl ozs/1,000 sq ft) after 5 to 8 weeks for extended <i>Poa annua</i> control.
		1.1 to 1.6	3.1 to 4.2	
<sup>1</sup> <b>DO NOT</b> exceed a maximum of 4.2 pints (2.1 quarts)/A or 1.6 fl ozs/1,000 sq ft product <b>per application</b> for use on residential and sod farm turfgrass. <b>DO NOT</b> exceed a maximum rate of 6.3 pints (3.1 quarts)/A or 2.3 fl ozs/1,000 sq ft product <b>per application</b> for use on golf course turfgrass, commercial, or other nonresidential turfgrass. <sup>2</sup> Residential is defined as turf in any residential situation as well as home lawns, schools, parks, and playgrounds. <sup>3</sup> <b>DO NOT</b> use on bentgrass or <i>Poa annua</i> greens or tees.				

## Weeds Controlled

**Pendulum AquaCap** will not control established weeds. If weeds should germinate before activation of herbicide, shallow cultivate to destroy existing weeds or, where practical, remove by hand. When cultivating for any reason, it should be shallow. **Pendulum AquaCap** may be used with herbicides registered for postemergence application (i.e. **Roundup® herbicide** or **Finale® herbicide**) for control of established weeds. **DO NOT** apply sprays containing **Roundup** or **Finale** over the top of desirable plants. A **Pendulum AquaCap** treatment may be followed by any registered herbicide to control weeds not listed on the **Pendulum AquaCap** label.

## Landscape and Grounds Maintenance

**Pendulum AquaCap** can be incorporated into landscape and grounds maintenance programs to provide extended preemergence control of most annual grasses and certain

broadleaf weeds. Areas to be treated, such as mulch beds, parking areas and roadsides, fencelines and borders, and around statuary or monuments, should be free of emerged weeds before application. To remove emerged weeds, either cultivate or tank mix **Pendulum AquaCap** with a postemergence product labeled for such use.

Not all ornamental species or cultivars of species can be tested for plant safety. Refer to the list of ornamental plant species found in this label (**Table 4. Tolerant Ornamental Species**). **Pendulum AquaCap** may be used on plant species not listed on this label; however, testing a small number plants at the specified rate and evaluating for suitability before a broad-use application is advised. Refer to **Table 3. Weed Control in All Nonturfgrass Sites** for application rates. Avoid unintentional contact of spray solution with stone, wood, or other porous surfaces because staining may occur. Rinse surfaces immediately using a heavy spray of water to avoid staining.

### Table 3. Weed Control in All Nonturfgrass Sites\*

For preemergence control of the weed species listed, apply **Pendulum® AquaCap herbicide** at the specified rates:

Length of Control (months)	Pendulum AquaCap (qts/A)	Required to Treat 1,000 sq ft (fl ozs)
Short term (2 to 4)	2.1	1.6
Long term (6 to 8)	4.2	3.2

\* For all turfgrass weed control rates, refer to **Table 2. Pendulum® AquaCap herbicide Residential, Golf Course, Commercial, and Other Nonresidential Turfgrass Uses for Preemergence Weed Control.**

For extended weed control, repeat applications of **Pendulum AquaCap** can be made.

### Ornamental Plantings and Tree Plantations including Noncropland Areas

Use **Pendulum AquaCap** for grounds maintenance in noncropland areas, for preemergence control of the weed species listed in and around established tree plantations for site preparation, and for maintenance of conifer and hardwood seedling nurseries and pulpwood and fiber farms. **Pendulum AquaCap** may be used for hardwood and conifer regeneration on conservation reserve program land. **Pendulum AquaCap** can also be used in Christmas trees and nonbearing fruit and nutcrops and vineyards established, or bulb and wildflower field plantings, in and around established ornamentals planted in noncropland areas such as highway rights-of-way and utility substations. Refer to **Table 3. Weed Control in All Nonturfgrass Sites** for application rates.

#### Applications at Planting or to Established Trees

When applying at planting, it is important to achieve slit closure to prevent **Pendulum AquaCap** from directly contacting the tree roots or being washed into the root zone via the open slit, or root stunting may occur. Refer to **Landscape and Ornamental Plantings Instructions and Restrictions** chart before application.

For postemergence weed control, tank mix combinations of **Pendulum AquaCap** plus **Segment® herbicide**, **Roundup® herbicide**, **Finale® herbicide**, or other labeled herbicides are recommended. Refer to approved labeling for species recommendations. Determine rates for tank mix compounds from the product labels of **Pendulum AquaCap** and partner herbicides before use. Use caution to prevent combination sprays from direct contact with desirable foliage or injury may result. **Pendulum AquaCap** plus diuron or simazine

combinations will broaden weed control spectrum; however, use of combinations may restrict **Pendulum AquaCap** use in sensitive areas. Refer to manufacturer's labels for specific use directions, precautions, and limitations before application and follow those that are most restrictive.

#### Ornamental Bulbs

**Pendulum AquaCap** may be applied for control of susceptible annual weeds in ornamental bulbs listed in the **Perennials** section in **Table 4. Tolerant Ornamental Species** (crocus, daffodil [narcissus], gladiolus, lily, tulip, etc.). Apply **Pendulum AquaCap** before, during, or after bulb emergence. If weeds have already germinated, add a labeled postemergence herbicide to control emerged weeds.

#### Wildflowers

**Pendulum AquaCap** may be applied for control of susceptible annual weeds in plantings of wildflowers listed in the **Perennials** section in **Table 4. Tolerant Ornamental Species**. The perennial species noted<sup>1</sup> (black-eyed Susan, California poppy, coreopsis, oxeye daisy, etc.) have been evaluated for plant tolerance to applications of **Pendulum AquaCap** at 4.2 pints (2.1 quarts) per acre. **Pendulum AquaCap** may be applied to established perennial wildflowers before emergence of weeds or wildflowers. For wildflowers being established from seed, apply **Pendulum AquaCap** no sooner than 4 weeks after wildflowers have emerged, but before weed germination. If weeds have already germinated, add a labeled postemergence product to control emerged weeds. Refer to all label restrictions before application.

Due to the diversity of species and varieties that exist in areas where wildflowers are grown, the response to **Pendulum AquaCap** may vary greatly. Careful testing on desirable species is recommended to determine if area-wide applications can be made.

#### Nonbearing Fruit and Nutcrops and Vineyards

**Pendulum AquaCap** may be applied for preemergence control of most annual grasses and certain broadleaf weeds on the following nonbearing crops:

Almond	Grape	Pistachio
Apple	Nectarine	Plum
Apricot	Olive	Prune
Cherry	Peach	Walnut, English
Citrus	Pear	
Fig	Pecan	

#### Noncropland

Use **Pendulum AquaCap** for preemergence control of most annual grasses and certain broadleaf weeds as they germinate on noncropland areas such as railroad, utility, highway, and pipeline rights-of-way; highway guardrails, delineators, and sign posts; utility substations, petroleum tank farms, pumping installations, fence rows, storage areas, windbreaks and shelterbelts.

## Industrial (Unimproved) Turf

**Pendulum® AquaCap herbicide** will provide pre-emergence control of the annual grasses and broadleaf weeds listed in **Table 1. Weeds Controlled** that might germinate in established grass in rights-of-way, roadsides, construction sites, parks, substations, or lots.

Apply before weeds germinate. A postemergence herbicide such as 2,4-D, **Drive® 75 DF herbicide**, **Drive® XLR8 herbicide**, **Segment® herbicide**, MSMA, or similar products may be tank mixed to control established weeds. Apply according to label instructions for the respective products and follow the most restrictive wording.

## Total Vegetation Control

**Pendulum AquaCap** may be tank mixed with **Arsenal® herbicide**, **Sahara® DG herbicide**, **Plateau® herbicide**, **Segment**, **Roundup PRO® herbicide**, **Karmex® herbicide**, **Finale® herbicide**, **Oust® herbicide**, diuron, glyphosate or other products to provide bareground or total vegetation control. **Pendulum AquaCap** can be used to provide greater plant selectivity in areas where such action may be desired. Such sites might have roots of landscape vegetation, ornamentals, or desirable trees encroaching into the treated zone. Refer to tank mix partner labels regarding effects on desirable plants. **DO NOT** tank mix with **Arsenal**, **Sahara DG**, or **Plateau herbicides** in California.

Applications may be made to existing weeds controlled by the partner herbicide. Determine rates from the product labels before use. Follow the most restrictive label instructions.

For kochia control, combinations of **Pendulum AquaCap** with **Arsenal** or diuron are recommended if control has been a problem for other herbicides.

## Landscape and Ornamental Plantings Instructions and Restrictions<sup>1</sup>

Site	Application Instructions and Restrictions
Landscape plantings <sup>2</sup>	<ol style="list-style-type: none"> <li><b>DO NOT</b> apply to newly transplanted ornamentals until plants have been watered and soil has been thoroughly packed and settled around roots.</li> <li>Apply as a directed or over-the-top spray.</li> <li>Use the lowest labeled rate when making applications to annuals. Repeat applications can be made for extended landscape weed control.</li> </ol>
Ornamental bulbs <sup>3</sup>	<ol style="list-style-type: none"> <li><b>Pendulum AquaCap</b> may be applied to bulb species listed on the label.</li> <li>Apply before bulb emergence.</li> </ol>

(continued)

## Landscape and Ornamental Plantings Instructions and Restrictions<sup>1</sup> (continued)

Site	Application Instructions and Restrictions
Wildflowers <sup>3</sup>	<ol style="list-style-type: none"> <li><b>Pendulum AquaCap</b> may be applied in plantings of wildflowers listed on the label. Refer to specific instructions for rate and plant tolerance.</li> <li>For wildflowers being established from seed, apply at 4 weeks after wildflowers have germinated, but before weed seed germination.</li> </ol>
<sup>1</sup> Plant only those desirable plant species listed on this label into soil treated the previous season with <b>Pendulum AquaCap</b> or injury may occur.	
<sup>2</sup> Before treating a large number of plants, spray a few plants and observe for 1 to 2 months for plant damage before full-scale application.	
<sup>3</sup> <b>DO NOT</b> treat plants grown for food or feed. <b>DO NOT</b> use treated plants for food or feed.	

## Weeds Controlled

**Pendulum AquaCap** will not control established weeds. If weeds germinate before herbicide activation, shallow cultivate to destroy existing weeds or, where practical, remove by hand. When cultivating for any reason, it should be shallow.

Use **Pendulum AquaCap** with herbicides registered for postemergence application (i.e. **Roundup® herbicide** or **Finale**) for control of established weeds. **DO NOT** apply sprays containing **Roundup** or **Finale** over the top of desirable plants. A **Pendulum AquaCap** treatment may be followed by any registered herbicide to control weeds not listed on the **Pendulum AquaCap** label.

The efficacy of **Pendulum AquaCap** will be improved if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. Erratic weed control may result if **Pendulum AquaCap** is not activated by rainfall or irrigation within 30 days.

## Commercial Ornamental Production

### Application Use Sites

**Pendulum AquaCap** can be used in and around field, liner, and container ornamental production.

**Pendulum AquaCap** sprays are safe around and over the top of the established plants listed in **Table 4. Tolerant Ornamental Species**. However, not all varieties or strains of the plant species listed have been tested. Refer to ornamental instructions and restrictions in this label before any application of **Pendulum AquaCap**. Unintentional consequences such as crop injury may result because of certain environmental or growing conditions, manner of use, or application. Therefore, before treating a large number of plants, spray a few plants and observe for plant damage before full-scale application.

## Application Instructions

**Pendulum® AquaCap herbicide** will not control established weeds. Therefore, areas to be treated should be free of established weeds at the time of treatment, or use **Pendulum AquaCap** with herbicides registered for post-emergence use in ornamentals and vegetation control sites. Consult the labels of those herbicides for suggested treatments, rates, and precautions or restrictions for use in these areas.

The efficacy of **Pendulum AquaCap** will improve if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. If **Pendulum AquaCap** is not activated by rainfall or irrigation within 30 days, erratic weed control may result.

Applied according to label directions and under normal growing conditions, **Pendulum AquaCap** or **Pendulum AquaCap** tank mix combinations will not cause crop injury. Overapplication can result in crop-stand loss, crop injury, or soil residues. Uneven application can decrease weed control or cause crop injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken seedlings and plants and increase the possibility of plant damage from **Pendulum AquaCap**.

## Production Ornamentals Instructions and Restrictions<sup>1</sup>

Site	Application Instructions and Restrictions
Newly transplanted field-grown nursery stock <sup>2,3</sup>	<ol style="list-style-type: none"> <li><b>DO NOT</b> make over-the-top applications at time of field transplanting. Use shielded sprayer until plantings have been established for one (1) year or more in the field.</li> <li><b>DO NOT</b> apply until transplants have been watered and soil has been thoroughly packed and settled around transplants. Take care to ensure there are no cracks in the soil where <b>Pendulum AquaCap</b> could come into contact with the roots.</li> <li><b>DO NOT</b> apply during bud swell, bud break, or at time of first flush of new growth.</li> <li>Direct sprays away from grafted or budded tissue on transplants at all times.</li> </ol>
Ornamental bulbs <sup>3</sup>	<ol style="list-style-type: none"> <li><b>Pendulum AquaCap</b> may be applied to bulb species listed on the label.</li> <li>Apply before bulb emergence.</li> </ol>

(continued)

## Production Ornamentals Instructions and Restrictions<sup>1</sup> (continued)

Site	Application Instructions and Restrictions
Newly transplanted container-grown nursery stock <sup>2,3</sup>	<ol style="list-style-type: none"> <li><b>DO NOT</b> apply until transplants have been watered and soil has been thoroughly packed and settled around transplants. Take care to ensure there are no cracks in the soil where <b>Pendulum AquaCap</b> could come into contact with the roots.</li> <li>For container-grown ornamentals, delay first application of the product to bareroot liners for two (2) weeks after transplanting.</li> <li><b>DO NOT</b> apply during bud swell, bud break, or at time of first flush of new growth.</li> <li>Direct sprays away from grafted or budded tissue on transplants at all times.</li> </ol>
Established container or field-grown nursery stock <sup>2,3</sup>	<ol style="list-style-type: none"> <li><b>DO NOT</b> apply during bud swell, bud break, or at time of first flush of new growth.</li> <li>Apply as a directed or over-the-top spray.</li> <li>If newly budded or grafted rootstock, apply with a shielded sprayer.</li> <li>Take care to ensure there are no cracks in the soil where <b>Pendulum AquaCap</b> could come into contact with the roots.</li> </ol>
Bareground for container placement	Apply to soil (including mulch, gravel, wood chips, or other permeable base); then water in. Replace containerized ornamentals onto pad.
Greenhouses, shadehouses, or other enclosed structures	<b>DO NOT</b> apply in greenhouses, shadehouses, or other enclosed structures.

<sup>1</sup> Plant only those desirable plant species listed on this label into soil treated the previous season with **Pendulum AquaCap** or injury may occur.

<sup>2</sup> Before treating a large number of plants, spray a few plants and observe for 1 to 2 months for plant damage before full-scale application.

<sup>3</sup> **DO NOT** treat plants grown for food or feed. **DO NOT** use treated plants for food or feed.

Refer to **Table 3. Weed Control in All Nonturfgrass Sites** for application rates.

## Ornamental Tank Mixes

Emerged weeds in ornamentals can be controlled using tank mixes containing **Segment® herbicide**, **Roundup® herbicide**, **Finale® herbicide**, **Ornamec® herbicide**, **Gallery® herbicide**, **Princep® herbicide**, and other similar products. **DO NOT** apply sprays containing **Roundup** or **Finale** over the top of ornamental plants.

Before tank mixing, use a simple jar test to ensure compatibility of herbicides.

Refer to manufacturer's labels for specific use directions, precautions, and limitations before tank mixing with **Pendulum® AquaCap herbicide**. Follow those that are most restrictive.

## Christmas Tree Plantations

Use **Pendulum AquaCap** in and around Christmas tree plantations. Apply **Pendulum AquaCap** at planting or to established trees. When applying at planting, it is important to achieve slit closure to prevent **Pendulum AquaCap** from directly contacting the tree roots or being washed into the root zone via the open slit or root stunting may occur.

For postemergence weed control, tank mix combinations of **Pendulum AquaCap** plus **Segment**, **Roundup**, **Finale**, or other labeled herbicides are recommended. Refer to approved labeling for species recommendations. Determine rates for tank mix combinations from the product labels of **Pendulum AquaCap** and partner herbicides before use. Use caution to prevent combination sprays from direct contact with desirable foliage or injury may result. **Pendulum AquaCap** plus diuron or simazine combinations will broaden weed control spectrum; however, use of combinations may restrict **Pendulum AquaCap** use in sensitive areas. Refer to manufacturer's labels for specific use directions, precautions, and limitations before application. Follow those that are most restrictive. Refer to **Table 3. Weed Control in All Nonturfgrass Sites** for **Pendulum AquaCap** application rates.

## Vegetation Control in Ornamental Production

Use **Pendulum AquaCap** for preemergence control of most annual grasses and certain broadleaf weeds as they germinate on noncropland areas such as sign posts, pumping installations, fence rows, storage areas, and windbreaks and shelterbelts. **Pendulum AquaCap** may be tank mixed with **Segment**, **Roundup PRO® herbicide**, **Karmex® herbicide**, **Finale**, diuron, glyphosate or other products to provide bareground or total vegetation control. **Pendulum AquaCap** can be used to provide greater plant selectivity in areas where such action may be desired. Such sites might have roots of landscape vegetation, ornamentals, or desirable trees encroaching into the treated zone. Refer to tank mix partner labels regarding effects on desirable plants. Applications may be made to existing weeds controlled by the partner herbicide. Determine rates from the product

labels before use. Follow the most restrictive label instructions. Refer to **Table 3. Weed Control in All Nonturfgrass Sites** for **Pendulum AquaCap** application rates.

## Weeds Controlled

**Pendulum AquaCap** will not control established weeds. If weeds germinate before herbicide activation, shallow cultivate to destroy existing weeds or, where practical, remove by hand. When cultivating for any reason, it should be shallow. **Pendulum AquaCap** may be used with herbicides registered for postemergence application (i.e. **Roundup** or **Finale**) for the control of established weeds. **DO NOT** apply sprays containing **Roundup** or **Finale** over the top of desirable plants. A **Pendulum AquaCap** treatment may be followed by any registered herbicide to control weeds not listed on the **Pendulum AquaCap** label.

The efficacy of **Pendulum AquaCap** will be improved if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. Erratic weed control may result if **Pendulum AquaCap** is not activated by rainfall or irrigation within 30 days.

**Pendulum AquaCap** may be used on plant species not listed on this label. Determine the suitability for such uses by treating a small number of such plants at the specified rate. Evaluate treated plants 1 to 2 months following treatment for possible injury.

**Pendulum AquaCap** sprays are safe around and over the top of the established plants listed in **Table 4. Tolerant Ornamental Species**. Refer to ornamentals instructions and restrictions before application. Refer to **Table 3. Weed Control in All Nonturfgrass Sites** for application rates.

**Table 4. Tolerant Ornamental Species**

Common Name	Scientific Name
<b>Bedding Plants</b>	
Ageratum	<i>Ageratum houstonianum</i>
Alyssum <sup>1</sup>	<i>Alyssum saxatile</i>
Anemone, poppy-flowered	<i>Anemone coronaria</i>
Artemesia	<i>Artemesia</i> spp.
Balloonflower	<i>Platycodon grandiflorum</i>
Begonia <sup>1</sup>	<i>Begonia</i> spp.
Cabbage, ornamental	<i>Brassica oleracea</i>
Caladium	<i>Caladium</i> spp.
Cast-iron plant	<i>Aspidistra elatior</i>
China aster <sup>1</sup>	<i>Callistephus chinensis</i>
Crocoshmia, monteбетria	<i>Crocoshmia x crocosmiiflora</i>
Dahlia <sup>1</sup>	<i>Dahlia</i> spp.
Dianthus	<i>Dianthus barbatus</i>

(continued)

**Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Bedding Plants</b> <i>(continued)</i>	
Dusty miller	<i>Senecio cineraria</i>
Gayfeather	<i>Liatris</i> spp.
Gazania, treasure flower	<i>Gazania rigens</i>
Gazania, trailing	<i>Gazania rigens leucolaena</i>
Gloxinia	<i>Gloxinia simningia</i>
Kale, ornamental	<i>Brassica napus</i>
Marigold, African	<i>Tagetes erecta</i>
Moss rose <sup>1</sup>	<i>Portulaca grandiflora</i>
Mum, garden	<i>Chrysanthemum</i> spp.
Periwinkle <sup>1</sup>	<i>Vinca major</i>
Periwinkle, rose	<i>Catharanthus roseus</i>
Petunia <sup>1</sup>	<i>Petunia</i> spp.
Plumosa cockscomb	<i>Celosia cristata</i>
Portulaca <sup>1</sup>	<i>Portulaca grandiflora</i>
Salvia <sup>1</sup>	<i>Salvia splendens</i>
Snapdragon	<i>Antirrhinum majus</i>
Statice <sup>1</sup>	<i>Limonium</i> spp.
Sweet William	<i>Dianthus barbatus</i>
Vinca <sup>1</sup>	<i>Vinca major</i>

<sup>1</sup> Application of **Pendulum® AquaCap herbicide** should not be made sooner than four weeks after transplanting for these annuals. Use the lower labeled rate.

**Ground Covers**

Ajuga	<i>Ajuga reptans</i>
Baby sun rose	<i>Aptenia cordifolia</i>
Beach strawberry	<i>Fragaria chiloensis</i>
Capeweed	<i>Arctotheca calendula</i>
Cinquefoil, spring	<i>Potentilla verna</i>
Coyotebrush, dwarf	<i>Baccharis pitularis</i>
Daisy, trailing African	<i>Osteospermum fruticosum</i>
Dymondia	<i>Dymondia margaretae</i>
Gazania	<i>Gazania splendens</i>
Iceplant, large leaf	<i>Carpobrotus edulis</i>
Ivy, English	<i>Hedera helix</i>
Ivy, geranium	<i>Pelargonium peltatum</i>
Jasmine, Asiatic	<i>Trachelospermum asiaticum</i>
Jasmine, primrose	<i>Jasminum mesnyi</i>
Jessamine, Carolina	<i>Gelsemium sempervirens</i>

*(continued)***Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Ground Covers</b> <i>(continued)</i>	
Manzanita, bearberry	<i>Arctostaphylos uva-ursi</i>
Miscanthus	<i>Miscanthus</i> spp.
Mondograss	<i>Ophiopogon japonica</i>
Morningglory	<i>Convolvulus</i> spp.
Myoporum	<i>Myoporum parvifolium</i>
Pachysandra	<i>Pachysandra terminalis</i>
Potentilla	<i>Potentilla fruticosa</i>
Red apple	<i>Aptenia cordifolia</i>
Rosemary	<i>Rosemarinus officinalis</i>
Rose-of-Sharon	<i>Hypericum calycinum</i>
St. Johnswort, creeping	<i>Hypericum calycinum</i>
Sand strawberry	<i>Fragaria chiloensis</i>
Sedum	<i>Sedum spurium</i>
Stonecrop	<i>Sedum spurium</i>
Verbena, Peruvian	<i>Verbena peruviana</i>
Vervain	<i>Verbena peruviana</i>
Vetch, crown	<i>Vicia sativa</i>
Vinca	<i>Vinca minor</i>
Wintercreeper	<i>Euonymus fortunei</i>

**Ornamental Grasses**

Beach grass	<i>Ammophila breviligulata</i>
Fescue, blue	<i>Festuca ovina</i>
Fescue, sheep	<i>Festuca ovina</i>
Fountain grass	<i>Pennisetum setaceum</i>
Pampas grass	<i>Cortaderia selloana</i>
Reed canary grass	<i>Phalaris arundinacea</i>
Reed, giant	<i>Arundo</i> spp.
Ribbon grass	<i>Phalaris arundinacea</i>
Tufted hair grass	<i>Deschampsia caespitosa</i>

**Perennials**

Acacia	<i>Acacia redolens</i>
Asparagus	<i>Asparagus</i> spp.
Aster, New York	<i>Aster novi-belgii</i>
Aster, Stokes	<i>Stokesia laevis</i>
Astilbe (False spirea)	<i>Astilbe</i> spp.
Avens	<i>Geum triflorum</i>
Baby's breath	<i>Gypsophila elegans</i>

*(continued)*

**Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Perennials</b> <i>(continued)</i>	
Baby's breath	<i>Gypsophila paniculata</i>
Beard-tongue	<i>Penstemon</i> spp.
Bellflower	<i>Campanula</i> spp.
Bellflower, willow	<i>Campanula persicifolia</i>
Bird of paradise	<i>Caesalpinia pulcherrima</i>
Black-eyed Susan <sup>1</sup>	<i>Rudbeckia hirta</i>
Blanket flower <sup>1</sup>	<i>Gaillardia aristata</i>
Blanket flower <sup>1</sup>	<i>Gaillardia x grandiflora</i>
Bleeding heart	<i>Dicentra spectabilis</i>
Butterfly weed	<i>Asclepias tuberosa</i>
California poppy <sup>1</sup>	<i>Eschscholzia californica</i>
Calla lily	<i>Zantedeschia aethiopica</i>
Canna, common garden	<i>Canna generalis</i> 'Lucifer'
Carex	<i>Carex</i> spp.
Chincherinchee	<i>Ornithogalum thyrsoides</i>
Clover, crimson <sup>1</sup>	<i>Trifolium incarnatum</i>
Columbine	<i>Aquilegia</i> 'McKana Giant'
Columbine	<i>Aquilegia x hybrida</i>
Coreopsis (Tickseed) <sup>1</sup>	<i>Coreopsis lanceolata</i>
Crinum lily	<i>Crinum</i> spp.
Crocus	<i>Crocus</i> spp.
Daffodil (Narcissus)	<i>Narcissus</i> spp.
Daylily	<i>Hemerocallis</i> spp.
Fairy duster	<i>Calliandra eriophylla</i>
Fern, asparagus	<i>Asparagus officinalis</i>
Fern, Boston	<i>Nephrolepis exaltata</i>
Fern, hay-scented	<i>Dennstaedtia punctilobula</i>
Fern, leatherleaf <sup>2</sup>	<i>Rumohra adiantiformis</i>
Fortnight lily	<i>Moraea</i> spp.
Foxglove	<i>Digitalis purpurea</i>
Freesia	<i>Freesia x hybrida</i>
Gaillardia	<i>Gaillardia pulchella</i>
Geum	<i>Geum</i> spp.
Gladiolus	<i>Gladiolus</i> spp.
Heather, dwarf	<i>Calluna vulgaris</i>
Hosta	<i>Hosta</i> spp.
Indian blanket <sup>1</sup>	<i>Gaillardia pulchella</i>
Iris, Japanese	<i>Iris kaempferi</i>

*(continued)***Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Perennials</b> <i>(continued)</i>	
Lantana, weeping	<i>Lantana montevidensis</i>
Leopard's bane	<i>Doronicum cordatum</i>
Lily	<i>Lilium</i> spp.
Liriope, big blue	<i>Liriope muscari</i>
Liriope, creeping	<i>Liriope spicata</i>
Liriope, variegated	<i>Liriope muscari</i>
Montbretia	<i>Crocsmia crocosmiiflora</i>
Moonbeam	<i>Coreopsis verticillata</i>
Mugwort, Western	<i>Artemesia ludoviciana</i>
Nightshade	<i>Solanum</i> spp.
Orchid, peacock	<i>Acidantha bicolor</i>
Oxeye daisy <sup>1</sup>	<i>Chrysanthemum leucanthemum</i>
Palm, areca	<i>Chyrsalidocarpus lutescens</i>
Palm, pygmy date	<i>Phoenix roebelenae</i>
Palm, Washington	<i>Washington robusta</i>
Peony, Chinese	<i>Paeonia lactiflora</i>
Purple coneflower <sup>1</sup>	<i>Echinacea purpurea</i>
Purple gay-feather	<i>Liatris pycnostachya</i>
Purple loosestrife	<i>Lythrum virgatum</i>
Rodgersia	<i>Rodgersia henricae</i>
Rosemary	<i>Rosmarinus officinalis</i>
Sedge	<i>Carex</i> spp.
Shasta daisy <sup>1</sup>	<i>Chrysanthemum x superbum</i>
Statice	<i>Limonium latifolia</i>
Statice, German	<i>Goniolimon tartaricum</i>
Sweet flag	<i>Acorus calamus</i>
Tickseed <sup>1</sup>	<i>Coreopsis lanceolata</i>
Texas bluebonnet	<i>Lupinus texensis</i>
Tulip	<i>Tulipa</i> spp.
Wonder flower	<i>Ornithogalum thyrsoides</i>
Yarrow <sup>1</sup>	<i>Achillea millefolium</i>
Zephyr lily	<i>Zephyranthes</i> spp.

<sup>1</sup> These plants have shown tolerance to **Pendulum® AquaCap herbicide** applications of 4.2 pints/A (2.1 quarts/A) in wildflower plantings established from seed.

<sup>2</sup> Applications of **Pendulum AquaCap** to immature ferns (during periods of new growth of fronds) may result in some injury.

*(continued)*

**Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Shrubs</b>	
Abelia, glossy	<i>Abelia grandiflora</i>
Alder, witch	<i>Fothergilla gardenii</i>
Aucuba, gold	<i>Aucuba japonica</i>
Azalea	<i>Rhododendron</i> spp.
Bamboo, heavenly	<i>Nandina domestica</i>
Barberry	<i>Berberis gladwynensis</i>
Barberry, Japanese	<i>Berberis thunbergii</i>
Blue indigo bush	<i>Dalea gregii</i>
Bottlebrush, lemon	<i>Callistemon citrinus</i>
Boxwood, common	<i>Buxus sempervirens</i>
Boxwood, Japanese	<i>Buxus microphylla</i>
Brittlebush	<i>Encelia farinosa</i>
Buttonbush	<i>Cephalanthus occidentalis</i>
Camellia	<i>Camellia japonica</i>
Cape jasmine	<i>Gardenia jasminoides</i>
Cassia, feathery	<i>Cassia artemisioides</i>
Cordyline	<i>Cordyline</i> spp.
Correa	<i>Correa</i> spp.
Cotoneaster	<i>Cotoneaster apiculatus</i>
Cotoneaster, bearberry	<i>Cotoneaster dammeri</i>
Cotoneaster, rock	<i>Cotoneaster horizontalis</i>
Cypress, Italian	<i>Cupressus sempervirens</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>
Deutzia, slender	<i>Deutzia gracilis</i>
Dogwood, red twig	<i>Cornus sericea</i>
Elaeagnus	<i>Elaeagnus ebbingei</i>
Escallonia	<i>Escallonia fradesii</i>
Euonymus	<i>Euonymus fortunei</i>
Euonymus, golden	<i>Euonymus japonica</i>
Euonymus, winged	<i>Euonymus alata</i>
Firethorn	<i>Pyracantha coccinea</i>
Forsythia, border	<i>Forsythia intermedia</i>
Fragrant olive	<i>Osmanthus fragrans</i>
Fuchsia, California	<i>Zauschneria californica</i>
Gardenia	<i>Gardenia jasminoides</i>
Hawthorne, Indian	<i>Raphiolepis indica</i>
Hibiscus	<i>Hibiscus syriacus</i>
Holly, Chinese	<i>Ilex cornuta</i>

*(continued)***Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Shrubs</b> <i>(continued)</i>	
Holly, Japanese	<i>Ilex crenata</i>
Holly, Fosters	<i>Ilex attenuata</i> 'Fosteri'
Holly, Savannah	<i>Ilex attenuata</i>
Holly, Yaupon	<i>Ilex vomitoria</i>
Honeysuckle, bush	<i>Diervilla lonicera</i>
Hopseed bush	<i>Dodonaea viscosa</i>
Hopbush	<i>Dodonaea viscosa</i>
Hydrangea	<i>Hydrangea macrophylla</i>
Juniper	<i>Juniperus</i> spp.
Juniper, Chinese	<i>Juniperus chinensis</i> v. <i>pfitzer</i>
Juniper, shore	<i>Juniperus conferta</i>
Juniper, trailing	<i>Juniperus horizontalis</i>
Laurel, cherry	<i>Prunus laurocerasus</i>
Laurel, mountain	<i>Kalmia latifolia</i>
Laurel, Otto Luyken	<i>Prunus laurocerasus</i>
Laurel, Schipka	<i>Prunus schipkanensis</i>
Laurustinus	<i>Viburnum tinus</i>
Lavender, English	<i>Lavandula angustifolia</i>
Leucothoe	<i>Leucothoe fontanesiana</i>
Leucothoe, coast	<i>Leucothoe axillaris</i>
Lilac, cut-leaf	<i>Syringa laciniata</i>
Lily-of-the-Nile	<i>Agapanthus africanus</i>
Mahonia	<i>Mahonia aquifolium</i>
Mock orange	<i>Pittosporum tobira</i>
Myrtle, compact	<i>Myrtus communis</i>
Myrtle, wax	<i>Myrica cerifera</i>
Nandina	<i>Nandina domestica</i>
Oleander	<i>Nerium oleander</i>
Oregon grape	<i>Mahonia aquifolium</i>
Osmanthus	<i>Osmanthus fragrans</i>
Palm, European fan	<i>Chamaerops humilis</i>
Palm, Mediterranean fan	<i>Chamaerops</i> spp.
Phlox, prickly	<i>Leptodactylon californicum</i>
Photinia, Fraser	<i>Photinia x fraseri</i>
Pieris, Japanese	<i>Pieris japonica</i>
Pine, Mugo	<i>Pinus mugo</i>
Plum, Natal	<i>Carissa grandiflora</i>
Privet, California	<i>Ligustrum ovalifolium</i>

*(continued)*

**Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Shrubs</b> <i>(continued)</i>	
Privet, glossy	<i>Ligustrum lucidum</i>
Privet, variegated	<i>Ligustrum sinensis</i>
Privet, waxleaf	<i>Ligustrum japonicum</i>
Pyracantha	<i>Pyracantha coccinea</i>
Quince, flowering	<i>Chaenomeles japonica</i>
Ranger, Texas	<i>Leucophyllum frutescens</i>
Redroot	<i>Ceanothus</i> spp.
Rhododendron	<i>Rhododendron</i> spp.
Robira	<i>Pittosporum tobira</i>
Rose	<i>Rosa</i> spp.
Spice plant	<i>Illicium parviflorum</i>
Spiraea	<i>Spiraea vanhouttei</i>
Spiraea, Anthony Waterer	<i>Spiraea x bumalda</i>
Spiraea, Japanese	<i>Spiraea japonica</i>
Sweet bay	<i>Laurus nobilis</i>
Trumpet bush	<i>Tecoma stans</i>
Verbena, lemon	<i>Aloysia triphylla</i>
Viburnum	<i>Viburnum suspensum</i>
Vitex	<i>Vitex</i> spp.
Weigela	<i>Weigela florida</i>
Wild lilac	<i>Ceanothus</i> spp.
Wisteria	<i>Wisteria</i> spp.
Xylosma	<i>Xylosma congestum</i>
Yellowbells	<i>Tecoma stans</i>
Yew <sup>1</sup>	<i>Taxus media</i>
Yew, Japanese <sup>1</sup>	<i>Taxus cuspidata</i>
Yew, Southern <sup>1</sup>	<i>Podocarpus macrophyllus</i>
Yucca, Adam's needle	<i>Yucca filamentosa</i>
Yucca, weeping	<i>Yucca pendula</i>

<sup>1</sup> Applications of **Pendulum® AquaCap herbicide** should not be made during spring growth or injury to terminals may occur.

**Trees**

Alder, European black	<i>Alnus glutinosa</i>
Apple	<i>Malus</i> spp.
Arborvitae, American	<i>Thuja occidentalis</i>
Arbutus	<i>Arbutus</i> spp.
Ash, red	<i>Fraxinus pennsylvanica</i>

*(continued)***Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Trees</b> <i>(continued)</i>	
Ash, white	<i>Fraxinus americana</i>
Aspen, bigtooth	<i>Populus grandidentata</i>
Aspen, quaking	<i>Populus tremuloides</i>
Basswood	<i>Tilia</i> spp.
Birch, European weeping	<i>Betula pendula</i>
Birch, river	<i>Betula nigra</i>
Buckeye, red	<i>Aesculus pavia</i>
Cedar, white	<i>Thuja occidentalis</i>
Chamaecyparis, Boulevard	<i>Chamaecyparis pisifera</i>
Cherry, black	<i>Prunus serotina</i>
Cherry, choke	<i>Prunus virginiana</i>
Cherry, Kwanzan	<i>Prunus serrulata</i>
Cherry, Nanking	<i>Prunus tomentosa</i>
Cottonwood	<i>Populus deltoides</i>
Crabapple	<i>Malus</i> spp.
Crape myrtle	<i>Lagerstroemia indica</i>
Cryptomeria, Japanese cedar	<i>Cryptomeria japonica</i>
Cypress, bald	<i>Taxodium distichum</i>
Cypress, Leyland	<i>Cupressocyparis leylandii</i>
Dogwood, flowering	<i>Cornus florida</i>
Dogwood, Korean	<i>Cornus kousa</i>
Dogwood, shrub	<i>Cornus</i> spp.
Dogwood, silky	<i>Cornus amomum</i>
Elm	<i>Ulmus japonica</i>
Elm, winged	<i>Ulmus alata</i>
Eucalyptus (Silver-dollar) tree	<i>Eucalyptus cinerea</i>
Fir, balsam	<i>Abies balsamae</i>
Fir, Douglas	<i>Pseudotsuga menziesii</i>
Fir, Fraser	<i>Abies fraseri</i>
Fir, white	<i>Abies concolor</i>
Franklinia	<i>Franklinia</i> spp.
Fringe tree	<i>Chlonenthus retusus</i>
Ginkgo	<i>Ginkgo biloba</i>
Gum, black	<i>Nyssa sylvatica</i>
Gum, sour	<i>Nyssa sylvatica</i>
Haw, black	<i>Viburnum prunifolium</i>

*(continued)*

**Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Trees (continued)</b>	
Hawthorn	<i>Crataegus</i> spp.
Hemlock, Canada	<i>Tsuga canadensis</i>
Hemlock, Eastern	<i>Tsuga canadensis</i>
Holly, American	<i>Ilex opaca</i>
Honeylocust	<i>Gleditsia triacanthos</i>
Lilac, common	<i>Syringa vulgaris</i>
Lilac, Japanese tree	<i>Syringa reticulata</i>
Linden	<i>Tilia</i> spp.
Magnolia, saucer	<i>Magnolia soulangiana</i>
Magnolia, Southern	<i>Magnolia grandiflora</i>
Magnolia, star	<i>Magnolia stellata</i>
Maidenhair tree	<i>Ginkgo biloba</i>
Maple, Japanese	<i>Acer palmatum</i>
Maple, Norway	<i>Acer platanoides</i>
Maple, red	<i>Acer rubrum</i>
Maple, sugar	<i>Acer saccharum</i>
Nannyberry, rusty	<i>Viburnum rufidulum</i>
Oak, chinquapin	<i>Quercus muehlenbergii</i>
Oak, live	<i>Quercus virginiana</i>
Oak, pin	<i>Quercus palustris</i>
Oak, red	<i>Quercus rubra</i>
Oak, swamp chestnut	<i>Quercus michauxii</i>
Oak, water	<i>Quercus nigra</i>
Oak, white	<i>Quercus alba</i>
Oak, willow	<i>Quercus phellos</i>
Olive	<i>Olea europaea</i>
Palm, date	<i>Phoenix</i> spp.
Palm, fan	<i>Washingtonia</i> spp.
Palm, pindo	<i>Butia</i> spp.
Palm, Washington	<i>Washingtonia</i> spp.
Peach	<i>Prunus persica</i>
Pear, Bradford	<i>Pyrus calleryana</i> 'Bradford'
Pecan	<i>Carya illinoensis</i>
Pine, Austrian	<i>Pinus nigra</i>
Pine, Italian stone	<i>Pinus pinea</i>
Pine, loblolly	<i>Pinus taeda</i>
Pine, Monterey	<i>Pinus radiata</i>
Pine, red	<i>Pinus resinosa</i>

*(continued)***Table 4. Tolerant Ornamental Species***(continued)*

Common Name	Scientific Name
<b>Trees (continued)</b>	
Pine, Scotch	<i>Pinus sylvestris</i>
Pine, Virginia	<i>Pinus virginiana</i>
Pine, white	<i>Pinus strobus</i>
Plum, purple leaf	<i>Prunus cerasifera</i>
Poplar, black	<i>Populus nigra</i>
Redcedar, Eastern	<i>Juniperus virginiana</i>
Redcedar, Western	<i>Thuja plicata</i>
Red ironbark	<i>Eucalyptus sideroxylon</i> 'Rosea'
Redwood, dawn	<i>Metasequoia glyptostroboides</i>
Sequoia, giant	<i>Sequoiadendron giganteum</i>
Serviceberry	<i>Amelanchier laevis</i>
Sourwood	<i>Oxydendrum arboreum</i>
Spruce, Colorado blue	<i>Picea pungens</i>
Spruce, dwarf Alberta	<i>Picea glauca</i> 'Albertiana'
Spruce, Norway	<i>Picea abies</i>
Spruce, white	<i>Picea glauca</i>
Sweetgum	<i>Liquidambar styraciflua</i>
Sycamore	<i>Platanus occidentalis</i>
Trachycarpus	<i>Trachycarpus</i> spp.
Tulip tree	<i>Liriodendron tulipifera</i>
Walnut, black	<i>Juglans nigra</i>
Willow, weeping	<i>Salix babylonica</i>
Yellowwood	<i>Cladrastis lutea</i>

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty**, which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

***Arsenal, Drive, Pendulum, Plateau, Sahara, and Segment** are registered trademarks of BASF.*

***Finale** is a registered trademark of Bayer.*

***Gallery** is a registered trademark of Dow Agrosciences LLC.*

***Karmex** is a registered trademark of ADAMA Group Company.*

***Oust** is a registered trademark of E. I. duPont de Nemours and Company.*

***Ornamec** and **Trimec** are registered trademarks of PBI Gordon Corp.*

***Princep** is a registered trademark of a Syngenta Group Company.*

***Roundup** and **Roundup PRO** are registered trademarks of Monsanto Technology LLC.*

***Three-Way** is a trademark of Lesco Technologies, LLC.*

© 2019 BASF Corporation  
All rights reserved.

000241-00416.20180702b.**NVA 2018-04-194-0050**  
Supersedes: NVA 2009-04-194-0050

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



We create chemistry

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29  
Version: 4.0

Page: 1/9  
(30255964/SDS\_CPA\_US/EN)

### 1. Product and Company Identification

Company  
BASF CORPORATION  
100 Park Avenue  
Florham Park, NJ 07932, USA

24 Hour Emergency Response Information  
CHEMTREC: 1-800-424-9300  
BASF HOTLINE: 1-800-832-HELP (4357)

---

Substance number:	000000171005
Molecular formula:	C13 H19 N3 O4
Chemical family:	aniline derivative
Synonyms:	pendimethalin

---

### 2. Hazards Identification

#### Emergency overview

**CAUTION:**  
Causes eye irritation.  
HARMFUL IF ABSORBED THROUGH SKIN.  
HARMFUL IF SWALLOWED.  
KEEP OUT OF REACH OF CHILDREN.  
KEEP OUT OF REACH OF DOMESTIC ANIMALS.  
Avoid contact with the skin, eyes and clothing.

See Product Label for additional precautionary statements.

State of matter: liquid  
Colour: yellow to brown  
Odour: faint odour, nutty

#### Potential health effects

##### **Primary routes of exposure:**

Routes of entry for solids and liquids include eye and skin contact, ingestion and inhalation. Routes of entry for gases include inhalation and eye contact. Skin contact may be a route of entry for liquified gases.

##### **Acute toxicity:**

Relatively nontoxic after single ingestion. Relatively nontoxic after short-term inhalation. Relatively nontoxic after short-term skin contact.

##### **Irritation / corrosion:**

May cause slight but temporary irritation to the eyes. May cause slight irritation to the skin.

##### **Sensitization:**

Skin sensitizing effects were not observed in animal studies.

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29  
Version: 4.0

Page: 2/9  
(30255964/SDS CPA US/EN)

### **Medical conditions aggravated by overexposure:**

Individuals with pre-existing diseases of the respiratory system, skin or eyes may have increased susceptibility to excessive exposures.

### **Signs and symptoms of overexposure:**

orange-red coloured urine caused by dye (not associated with methemoglobinemia)

### **Potential environmental effects**

#### **Aquatic toxicity:**

Very toxic (acute effect) to aquatic organisms.

#### **Terrestrial toxicity:**

Acutely harmful to terrestrial organisms.

## 3. Composition / Information on Ingredients

<u>CAS Number</u>	<u>Content (W/W)</u>	<u>Chemical name</u>
40487-42-1	38.7 %	pendimethalin
	61.3 %	Proprietary ingredients

## 4. First-Aid Measures

### **General advice:**

First aid providers should wear personal protective equipment to prevent exposure. Remove contaminated clothing. Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or physician for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

### **If inhaled:**

Remove the affected individual into fresh air and keep the person calm.

### **If on skin:**

Rinse skin immediately with plenty of water for 15 - 20 minutes.

### **If in eyes:**

Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing.

### **If swallowed:**

Never induce vomiting or give anything by mouth if the victim is unconscious or having convulsions. Do not induce vomiting. Have person sip a glass of water if able to swallow.

### **Note to physician**

Antidote: No known specific antidote.  
Treatment: Treat symptomatically.

## 5. Fire-Fighting Measures

Flash point:	> 230 °F	
Autoignition:	354 °C	(DIN EN 14522)
Lower explosion limit:		not determined
Upper explosion limit:		not determined
Flammability:	not highly flammable	

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29  
Version: 4.0

Page: 3/9  
(30255964/SDS CPA US/EN)

### **Suitable extinguishing media:**

foam, dry powder, carbon dioxide, water spray

### **Hazards during fire-fighting:**

carbon monoxide, carbon dioxide, nitrogen dioxide, nitrogen oxide, Hydrocarbons,  
If product is heated above decomposition temperature, toxic vapours will be released. The substances/groups of substances mentioned can be released in case of fire.

### **Protective equipment for fire-fighting:**

Firefighters should be equipped with self-contained breathing apparatus and turn-out gear.

### **Further information:**

Evacuate area of all unnecessary personnel. Contain contaminated water/firefighting water. Do not allow to enter drains or waterways.

---

## 6. Accidental release measures

### **Personal precautions:**

Take appropriate protective measures. Clear area. Shut off source of leak only under safe conditions. Extinguish sources of ignition nearby and downwind. Ensure adequate ventilation. Wear suitable personal protective clothing and equipment.

### **Environmental precautions:**

Do not discharge into the subsoil/soil. Do not discharge into drains/surface waters/groundwater. Contain contaminated water/firefighting water.

### **Cleanup:**

Dike spillage. Pick up with suitable absorbent material. Place into suitable containers for reuse or disposal in a licensed facility. Spilled substance/product should be recovered and applied according to label rates whenever possible. If application of spilled substance/product is not possible, then spills should be contained, solidified, and placed in suitable containers for disposal. After decontamination, spill area can be washed with water. Collect wash water for approved disposal.

---

## 7. Handling and Storage

### Handling

#### **General advice:**

RECOMMENDATIONS ARE FOR MANUFACTURING, COMMERCIAL BLENDING, AND PACKAGING WORKERS. PESTICIDE APPLICATORS & WORKERS must refer to the Product Label and Directions for Use attached to the product for Agricultural Use Requirements in accordance with the EPA Worker Protection Standard 40 CFR part 170. Ensure adequate ventilation. Provide good ventilation of working area (local exhaust ventilation if necessary). Keep away from sources of ignition - No smoking. Keep container tightly sealed. Protect contents from the effects of light. Protect against heat. Protect from air. Handle and open container with care. Do not open until ready to use. Once container is opened, content should be used as soon as possible. Avoid aerosol formation. Avoid dust formation. Provide means for controlling leaks and spills. Do not return residues to the storage containers. Follow label warnings even after container is emptied. The substance/product may be handled only by appropriately trained personnel. Avoid all direct contact with the substance/product. Avoid contact with the skin, eyes and clothing. Avoid inhalation of dusts/mists/vapours. Wear suitable personal protective clothing and equipment.

#### **Protection against fire and explosion:**

The relevant fire protection measures should be noted. Fire extinguishers should be kept handy. Avoid all sources of ignition: heat, sparks, open flame. Sources of ignition should be kept well clear. Avoid extreme heat. Keep away from oxidizable substances. Electrical equipment should conform to national electric code. Ground all transfer equipment properly to prevent electrostatic discharge. Electrostatic discharge may cause ignition.

### Storage

#### **General advice:**

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29  
Version: 4.0

Page: 4/9  
(30255964/SDS\_CPA\_US/EN)

Keep only in the original container in a cool, dry, well-ventilated place away from ignition sources, heat or flame. Protect containers from physical damage. Protect against contamination. The authority permits and storage regulations must be observed.

### **Storage incompatibility:**

General advice: Segregate from incompatible substances. Segregate from foods and animal feeds. Segregate from textiles and similar materials.

### **Temperature tolerance**

Protect from temperatures below: 0 °C

Changes in the properties of the product may occur if substance/product is stored below indicated temperature for extended periods of time.

Protect from temperatures above: 40 °C

Changes in the properties of the product may occur if substance/product is stored above indicated temperature for extended periods of time.

---

## 8. Exposure Controls and Personal Protection

**Users of a pesticidal product should refer to the product label for personal protective equipment requirements.**

### **Advice on system design:**

Whenever possible, engineering controls should be used to minimize the need for personal protective equipment.

### **Personal protective equipment**

#### **RECOMMENDATIONS FOR MANUFACTURING, COMMERCIAL BLENDING, AND PACKAGING WORKERS:**

##### **Respiratory protection:**

Wear respiratory protection if ventilation is inadequate. Wear a NIOSH-certified (or equivalent) TC23C Chemical/Mechanical type filter system to remove a combination of particles, gas and vapours. For situations where the airborne concentrations may exceed the level for which an air purifying respirator is effective, or where the levels are unknown or Immediately Dangerous to Life or Health (IDLH), use NIOSH-certified full facepiece pressure demand self-contained breathing apparatus (SCBA) or a full facepiece pressure demand supplied-air respirator (SAR) with escape provisions.

##### **Hand protection:**

Chemical resistant protective gloves, Protective glove selection must be based on the user's assessment of the workplace hazards.

##### **Eye protection:**

Safety glasses with side-shields. Tightly fitting safety goggles (chemical goggles). Wear face shield if splashing hazard exists.

##### **Body protection:**

Body protection must be chosen depending on activity and possible exposure, e.g. head protection, apron, protective boots, chemical-protection suit.

##### **General safety and hygiene measures:**

Wear long sleeved work shirt and long work pants in addition to other stated personal protective equipment. Work place should be equipped with a shower and an eye wash. Handle in accordance with good industrial hygiene and safety practice. Personal protective equipment should be decontaminated prior to reuse. Gloves must be inspected regularly and prior to each use. Replace if necessary (e.g. pinhole leaks). Take off immediately all contaminated clothing. Store work clothing separately. Hands and/or face should be washed before breaks and at the end of the shift. No eating, drinking, smoking or tobacco use at the place of work. Keep away from food, drink and animal feeding stuffs.

---

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29  
Version: 4.0

Page: 5/9  
(30255964/SDS CPA US/EN)

### 9. Physical and Chemical Properties

Form:	microencapsulated, suspension	
Odour:	faint odour, nutty	
Colour:	yellow to brown	
pH value:	approx. 8 - 10	( 1 %(m), 21 °C)
Melting point:	approx. 0 °C	Information applies to the solvent.
Density:	approx. 9.79 lb/USg	( 68 °F)
Vapour density:		not determined
Partitioning coefficient n-octanol/water (log Pow):		not applicable
Viscosity, dynamic:	128 mPa.s	( 20 °C) (OECD 114)
Solubility in water:		dispersible
Molar mass:	281.31 g/mol	

### 10. Stability and Reactivity

#### Conditions to avoid:

Avoid all sources of ignition: heat, sparks, open flame. Avoid prolonged storage. Avoid electro-static discharge. Avoid contamination. Avoid prolonged exposure to extreme heat. Avoid extreme temperatures.

#### Substances to avoid:

strong acids, strong bases, strong oxidizing agents

#### Hazardous reactions:

The product is chemically stable.

Hazardous polymerization will not occur. No hazardous reactions if stored and handled as prescribed/indicated.

#### Decomposition products:

No hazardous decomposition products if stored and handled as prescribed/indicated., Prolonged thermal loading can result in products of degradation being given off.

#### Thermal decomposition:

Possible thermal decomposition products:

carbon monoxide, carbon dioxide, nitrogen dioxide, nitrogen oxide, Hydrocarbons

Stable at ambient temperature. If product is heated above decomposition temperature toxic vapours may be released.

#### Oxidizing properties:

not fire-propagating

### 11. Toxicological information

#### Acute toxicity

##### Oral:

Type of value: LD50

Species: rat (male/female)

Value: > 5,000 mg/kg (OECD Guideline 401)

##### Inhalation:

Type of value: LC50

Species: rat (male/female)

Value: > 5.23 mg/l (OECD Guideline 403)

Exposure time: 4 h

An aerosol was tested.

No mortality was observed.

##### Dermal:

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29

Version: 4.0

Page: 6/9

(30255964/SDS CPA\_US/EN)

Type of value: LD50  
Species: rat (male/female)  
Value: > 5,000 mg/kg (OECD Guideline 402)

### Irritation / corrosion

#### Skin:

Species: rabbit  
Result: mildly irritating

#### Eye:

Species: rabbit  
Result: mildly irritating

#### Sensitization:

modified Buehler test  
Species: guinea pig  
Result: Non-sensitizing.  
Method: OECD Guideline 406

### Repeated dose toxicity

*Information on: pendimethalin*

*Assessment of repeated dose toxicity:*

*No substance-specific organotoxicity was observed after repeated administration to animals. Adaptive effects were observed after repeated exposure in animal studies.*

*Information on: Diphenylmethane-4,4'-diisocyanate (MDI)*

*Assessment of repeated dose toxicity:*

*The substance may cause damage to the olfactory epithelium after repeated inhalation. These effects are not relevant to humans at occupational levels of exposure.*

*Information on: Methylenediphenyl diisocyanate*

*Assessment of repeated dose toxicity:*

*The substance may cause damage to the olfactory epithelium after repeated inhalation. These effects are not relevant to humans at occupational levels of exposure.*

### Carcinogenicity

*Information on: pendimethalin*

*In long-term studies in rats the substance induced thyroid tumors. The effect is caused by an animal specific mechanism that has no human counter part. In long-term studies in mice in which the substance was given by feed, a carcinogenic effect was not observed.*

*Information on: Diphenylmethane-4,4'-diisocyanate (MDI)*

*A carcinogenic potential cannot be excluded after prolonged exposure to severely irritating concentrations. These effects are not relevant to humans at occupational levels of exposure.*

*Information on: Methylenediphenyl diisocyanate*

*A carcinogenic potential cannot be excluded after prolonged exposure to severely irritating concentrations. These effects are not relevant to humans at occupational levels of exposure.*

### Development:

*Information on: Diphenylmethane-4,4'-diisocyanate (MDI)*

*The substance did not cause malformations in animal studies; however, toxicity to development was observed at high doses that were toxic to the parental animals.*

*Information on: Methylenediphenyl diisocyanate*

*The substance did not cause malformations in animal studies; however, toxicity to development was observed at high doses that were toxic to the parental animals.*

### Experiences in humans:

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29

Version: 4.0

Page: 7/9

(30255964/SDS CPA US/EN)

Pendimethalin is a strongly orange-red compound - virtually an aniline dye. Cases have been described of orange-yellow colouration of urine following heavy exposure of workers to the dust of pendimethalin. Despite its structure as both a nitro-compound and aromatic amine, exposure to pendimethalin is NOT associated with methemoglobinemia.

### Other Information:

Misuse can be harmful to health.

---

## 12. Ecological Information

### Fish

Acute:

OECD Guideline 203 static

Oncorhynchus mykiss/LC50 (96 h): 20.36 mg/l

### Aquatic invertebrates

Acute:

OECD Guideline 202, part 1 static

Daphnia magna/EC50 (48 h): > 100 mg/l

### Aquatic plants

Toxicity to aquatic plants:

OECD Guideline 201 green algae/EC50 (72 h): 1.49 mg/l

### Non-Mammals

*Information on: pendimethalin*

*Other terrestrial non-mammals:*

*mallard duck/LD50: 1,421 mg/kg*

*Acutely harmful to terrestrial organisms.*

*Honey bee/LD50: 49.8 ug/bee*

*Acutely harmful to terrestrial organisms.*

### Environmental mobility:

*Information on: pendimethalin*

*Assessment transport between environmental compartments:*

*Following exposure to soil, adsorption to solid soil particles is probable, therefore contamination of groundwater is not expected.*

### Other adverse effects:

The ecological data given are those of the active ingredient. Do not release untreated into natural waters.

---

## 13. Disposal considerations

### Waste disposal of substance:

Pesticide wastes are regulated. Improper disposal of excess pesticide, spray mix or rinsate is a violation of federal law. If pesticide wastes cannot be disposed of according to label instructions, contact the State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29  
Version: 4.0

Page: 8/9  
(30255964/SDS CPA US/EN)

### Container disposal:

Rinse thoroughly at least three times (triple rinse) in accordance with EPA recommendations. Consult state or local disposal authorities for approved alternative procedures such as container recycling. Recommend crushing, puncturing or other means to prevent unauthorized use of used containers.

### RCRA: D028

The waste codes are manufacturer's recommendations based on the designated use of the product. Other use and special waste disposal treatment on customer's location may require different waste-code assignments.

---

## 14. Transport Information

### Land transport

USDOT

Not classified as a dangerous good under transport regulations

### Sea transport

IMDG

Hazard class: 9  
Packing group: III  
ID number: UN 3082  
Hazard label: 9, EHSM  
Marine pollutant: YES  
Proper shipping name: ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.  
(contains PENDIMETHALIN)

### Air transport

IATA/ICAO

Hazard class: 9  
Packing group: III  
ID number: UN 3082  
Hazard label: 9, EHSM  
Proper shipping name: ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.  
(contains PENDIMETHALIN)

### Further information

DOT: This product is regulated if the amount in a single receptacle exceeds the Reportable Quantity (RQ). Please refer to Section 15 of this MSDS for the RQ for this product.

---

## 15. Regulatory Information

### Federal Regulations

#### Registration status:

Crop Protection TSCA, US released / listed

Chemical TSCA, US blocked / not listed

OSHA hazard category: Skin and/or eye irritant; Chronic target organ effects reported

EPCRA 311/312 (Hazard categories): Acute; Chronic

EPCRA 313:

# Safety Data Sheet

## PENDULUM AQUACAP HERBICIDE

Revision date : 2012/11/29

Page: 9/9

Version: 4.0

(30255964/SDS\_CPA\_US/EN)

**CAS Number**

40487-42-1

**Chemical name**

pendimethalin

---

### 16. Other Information

**Refer to product label for EPA registration number.**

Recommended use: herbicide

**NFPA Hazard codes:**

Health : 1      Fire: 1      Reactivity: 1      Special:

We support worldwide Responsible Care® initiatives. We value the health and safety of our employees, customers, suppliers and neighbors, and the protection of the environment. Our commitment to Responsible Care is integral to conducting our business and operating our facilities in a safe and environmentally responsible fashion, supporting our customers and suppliers in ensuring the safe and environmentally sound handling of our products, and minimizing the impact of our operations on society and the environment during production, storage, transport, use and disposal of our products.

---

**MSDS Prepared by:**

BASF NA Product Regulations

msds@basf.com

MSDS Prepared on: 2012/11/29

IMPORTANT: WHILE THE DESCRIPTIONS, DESIGNS, DATA AND INFORMATION CONTAINED HEREIN ARE PRESENTED IN GOOD FAITH AND BELIEVED TO BE ACCURATE , IT IS PROVIDED FOR YOUR GUIDANCE ONLY. BECAUSE MANY FACTORS MAY AFFECT PROCESSING OR APPLICATION/USE, WE RECOMMEND THAT YOU MAKE TESTS TO DETERMINE THE SUITABILITY OF A PRODUCT FOR YOUR PARTICULAR PURPOSE PRIOR TO USE. NO WARRANTIES OF ANY KIND, EITHER EXPRESSED OR IMPLIED, INCLUDING WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE MADE REGARDING PRODUCTS DESCRIBED OR DESIGNS, DATA OR INFORMATION SET FORTH, OR THAT THE PRODUCTS, DESIGNS, DATA OR INFORMATION MAY BE USED WITHOUT INFRINGING THE INTELLECTUAL PROPERTY RIGHTS OF OTHERS. IN NO CASE SHALL THE DESCRIPTIONS, INFORMATION, DATA OR DESIGNS PROVIDED BE CONSIDERED A PART OF OUR TERMS AND CONDITIONS OF SALE. FURTHER, YOU EXPRESSLY UNDERSTAND AND AGREE THAT THE DESCRIPTIONS, DESIGNS, DATA, AND INFORMATION FURNISHED BY OUR COMPANY HEREUNDER ARE GIVEN GRATIS AND WE ASSUME NO OBLIGATION OR LIABILITY FOR THE DESCRIPTION, DESIGNS, DATA AND INFORMATION GIVEN OR RESULTS OBTAINED, ALL SUCH BEING GIVEN AND ACCEPTED AT YOUR RISK.  
END OF DATA SHEET