



NOTICE OF PESTICIDE APPLICATION

DATE OF APPLICATION: Week of 11/18/2024

LOCATION:

1. Trails Linear Park - Between Spring Gate Ln. & Snow Trail

PRODUCT & MANUFACTURER INFORMATION:

Product: Safari 20 SG Insecticide

Manufacturer: Valent U.S.A. LLC

EPA Reg. No. 86203-11-59639

Active ingredients(S): Dinotefurean, [N-methyl-N'-nitro-N''-(tetrahydro-3-furanyl)methyl] guanidine]..20%

Precautionary statement: CAUTION! Hazards to humans and domestic animals. Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Causes moderate eye irritation. Remove and wash contaminated clothing before reuse.


REASON FOR APPLICATION: treating Honey Mesquite Trees for Boring Insects

*Attached is the label and SDS sheet

Safari®

20 SG INSECTICIDE

GROUP 4A INSECTICIDE



FOR FOLIAR AND SYSTEMIC INSECT CONTROL IN ORNAMENTAL PLANTS AND VEGETABLE TRANSPLANTS IN ENCLOSED STRUCTURES. FOR GREENHOUSE, NURSERY, INTERIOR PLANTSCAPE, OUTDOOR LANDSCAPE, GREENHOUSE TOMATO AND FORESTRY USE ONLY.

Active Ingredient:
 Dinotefuran, [N-methyl-N'-nitro-N''-(tetrahydro-3-furanyl)methyl]guanidine].. 20%
 Other Ingredients..... 80%
 Total 100%
 EPA Reg. No. 86203-11-59639
 EPA Est. 67545-AZ-1

KEEP OUT OF REACH OF CHILDREN CAUTION
SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS
NET CONTENTS 3 POUNDS



(01) 0 06 53204 07571 0 Form 1510-I

FIRST AID

- If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
- If swallowed:** Call poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by the poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
- If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
- If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Causes moderate eye irritation. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made out of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or viton \geq 14 mils.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not dispose equipment washwaters or rinsate into a natural drain or water body.

This product is toxic to honey bees. The persistence of residues and potential residual toxicity of dinotefuran in nectar and pollen suggests the possibility of chronic toxic risk to honey bee larvae and the eventual instability of the hive.

- This product is toxic to bees exposed to residues for more than 38 hours following treatment.
- Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

Dinotefuran and its degradate, MNG, have the properties and characteristics associated with chemicals detected in groundwater. The high water solubility of dinotefuran, and its degradate, MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.



PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

SPRAY DRIFT ADVISORY

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered for sale, use or consumption.

PROTECTION OF POLLINATORS
APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

(continued)

PROTECTION OF POLLINATORS (continued)

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>. Pesticide incidents (for example, bee kills) should immediately be reported to the State/Tribal lead agency. For contact information for your State, go to: www.aapco.org. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

FOR COMMERCIALLY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS



- Do not apply this product while bees are foraging.
- This product is toxic to bees exposed to residue for more than 38 hours following treatment.
- Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

NON-AGRICULTURAL USES



Do not apply Safari® 20 SG Insecticide while bees are foraging. Do not apply Safari 20 SG Insecticide to plants that are flowering. Only apply after all flower petals have fallen off.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural insecticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

EXCEPTION: If product is drenched or soil-injected, workers may enter the area at any time if there will be no contact with anything that has been treated.

(continued)

(continued)

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves (made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or viton \geq 14 mils)

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow others to enter treated areas until sprays have dried.

APPLICATION INFORMATION

- Applications of *Safari 20 SG* Insecticide in residential areas may be made by commercially licensed applicators.

Application to Ornamental Plants (including Forestry):

- *Safari 20 SG* Insecticide can be applied as a foliar spray, a broadcast spray, a soil drench, soil injection and via chemigation for insect control in ornamental plants in greenhouses, nurseries, outdoor landscapes and interior plantscapes.
- *Safari 20 SG* Insecticide is a systemic product and will be taken up by the root system and trans-

located upward throughout the plant. When applied as a foliar spray, the product offers translaminar and locally systemic control of foliar pests.

- When applied to the soil, *Safari 20 SG* Insecticide will be translocated more quickly in herbaceous plants than in woody shrubs and trees. Speed of insect control will range from as little as one day for small herbaceous plants in containers, to several weeks in large trees growing in the landscape.
- Do not apply more than a total of 2.7 lbs of product (0.54 lb active ingredient) per acre per year for all application types.
- **Do not apply this product, by any application method, to linden, basswood or other *Tilia* species.**
- **Use on *Myoporum laetum* is not recommended.**

Application to Vegetable Transplants:

- *Safari 20 SG* Insecticide can be applied as a foliar spray or a broadcast spray for insect control in vegetable transplants.
- Do not apply more than 1.34 lbs (0.268 lbs ai) per acre of nursery per year.

MIXING INSTRUCTIONS:

Safari 20 SG Insecticide Alone: Add half of the required amount of water to the mix tank. With the agitator running, add the desired amount of *Safari 20 SG* Insecticide to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after *Safari 20 SG* Insecticide has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Safari 20 SG Insecticide + Tank Mixtures: Add half of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, emulsifiable concentrates, and surfactants/adjuvants.

Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

NOTE: When using *Safari 20 SG* Insecticide in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including *Safari 20 SG* Insecticide. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using *Safari 20 SG* Insecticide in a tank mixture, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Do not exceed label dosage rate. Do not mix this product with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Volumes, weights, and conversions

Use the following equivalent, unpacked volumetric weights

1 tablespoon = 0.22 oz = approximately 1/4 oz;
2.5 tablespoons = 0.55 oz = approximately 1/2 oz;
3.5 tablespoons = 0.77 oz = approximately 3/4 oz;
4.5 tablespoons = 0.99 oz = approximately 1 oz;
28 grams = 1.0 oz
2.7 lb = 43.2 oz

Compatibility

IMPORTANT: The safety of all potential tank mixes has not been tested on all crops. Before applying any tank mixture not specifically listed on this label, confirm the safety to the target crop.

Safari 20 SG Insecticide is compatible with most commonly used pesticides, crop oils, adjuvants, and nutritional sprays. However, since it is not possible to test all possible mixtures, pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with *Safari 20 SG* Insecticide. To determine the physical compatibility of *Safari 20 SG* Insecticide with other products, use a jar test, as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for additional required ingredients to the spray tank.

RESISTANCE MANAGEMENT

Safari 20 SG Insecticide contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly in the same crop or in successive years as the primary method of control for a targeted species. This may result in partial or total loss of control of those species by *Safari 20 SG* Insecticide or other Group 4A insecticides.

To delay the development of insecticide resistance in greenhouse, nursery and interiorscape use sites, strongly consider the following guidelines:

- Do not apply *Safari 20 SG* Insecticide or other Group 4A insecticides to consecutive generations of the same insect pest species.
- Do not drench soil media with *Safari 20 SG* Insecticide or other Group 4A insecticides more than one time per crop cycle or three months, whichever is shorter.
- Do not make more than two foliar or broadcast sprays of *Safari 20 SG* Insecticide or other Group 4A insecticides to a single crop during a two-month period.
- Do not make more than one soil drench and one foliar or broadcast spray with *Safari 20 SG* Insecticide or other Group 4A insecticides during a two-month period.
- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturers for insecticide resistance management and/or IPM guidelines for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Valent U.S.A. LLC, at toll free number: 1-800-898-2536.

APPLICATION PROCEDURES AND SPRAY EQUIPMENT

Ground Application: Select spray nozzles that will provide accurate and uniform spray deposition. Use spray nozzles that provide medium-sized droplets and reduce drift. To help insure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult nozzle manufacturers and/or State Extension Service specialists.

Apply *Safari 20 SG* Insecticide using sufficient water volume to provide thorough and uniform coverage. In situations where a dense canopy exists and/or pest pressure is high, use greater water volumes. The use of a spray adjuvant may improve spray coverage. Do not apply under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Applications to ornamental plants, forestry and vegetable transplants: *Safari 20 SG* Insecticide can be applied using many different types of application equipment. Apply in sufficient water to ensure good coverage of ornamental plants. Tank mixing with a surfactant will produce better coverage when making applications to plants with hard to wet foliage such as holly or pine. If concentrate or mist type spray equipment is used, apply the same amount of product on the sprayed area as would be used in a dilute solution. To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. Applications can be made to foliage or as a soil drench.

RESTRICTIONS

- With the exception of non-livestock animals, do not graze treated areas or use clippings from treated areas for feed or forage.
- Prevent runoff or puddling of irrigation water following application.
- Keep children and pets off treated areas until spray has dried.
- Do not apply to areas that are water logged or saturated, or frozen, which will not allow penetration into the root zone of the plant.

APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION):

Safari 20 SG Insecticide may be applied by injection into an irrigation system, either alone or in combination with other pesticides or chemicals that are registered for application through irrigation systems. Dilution ratios are normally 1:100 to 1:200, depending on the system. Apply this product only through microirrigation (individual spaghetti tube), drip irrigation, overhead irrigation, or motorized calibrated irrigation equipment (Ornamentals). Do not apply through any other type of irrigation system. Lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make adjustments when necessary.

Using Water from Public Water Systems:**DO NOT APPLY SAFARI 20 SG INSECTICIDE THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM.**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year. *Safari 20 SG* Insecticide may be applied through irrigation systems which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements:

Operating Instructions for Irrigation Systems:

1. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
2. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Do not apply when wind speed favors drift beyond the area intended.

Calibration and Application Instructions:

Apply *Safari 20 SG* Insecticide under the schedule specified in the specific use instructions, not according to the irrigation schedule unless the events coincide. In general, set the equipment to apply the minimum amount of water per acre. Run the system at 86-90% of the manufacturer's maximum rated travel speed.

The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Check with State and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

MINIMIZING SPRAY DRIFT

As with all crop protection products, it is important to minimize off-target movement. Do not allow spray to drift onto adjacent land, crops, or aquatic areas. To minimize spray drift:

1. Make applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 10 mph. Do not apply when wind gusts approach 10 mph.
2. Risk of exposure to sensitive aquatic areas can be reduced by not applying when wind direction is toward the aquatic area.
3. Do not cultivate or plant crops within 25 feet of the aquatic area as to allow growth of a vegetative filter strip.
4. Do not make applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with increased height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

5. Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by not using excessive spray boom pressure.
6. Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage. Do not apply more than 10 ft above the crop canopy.
7. For aerial applications, mount spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use minimum practical boom length and do not use boom that exceeds 75% of wing span or rotor diameter.

Air Assisted (Air Blast) Tree and Vine Sprayers (Ornamentals Only):

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift:

1. Adjust deflectors and aiming devices so that spray is only directed into the canopy.
2. Block off upward pointed nozzles when there is no overhanging canopy.
3. Use only enough air volume to penetrate the canopy and provide good coverage. Use a minimum of 50 gallons finished spray per acre.
4. Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.

**VEGETABLE TRANSPLANTS (IN ENCLOSED STRUCTURES)
FOLIAR OR BROADCAST SPRAY APPLICATION**

For foliar insect control on vegetable transplants grown in enclosed structures.

Crops	Pests	Product Rate (By Weight)	Remarks
Cucurbits (Transplants only)	Aphids	3.5-7.0 oz per 100 gal	Do not make more than one application per crop.
Cantaloupe	Leafminers	7 - 14 oz per Acre	Apply only to cucurbits and brassica being grown as transplants and before transplants are sold.
Cucumber	Mealybugs	0.16-0.32 oz per 1,000 sq ft	100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Melons	Thrips (suppression)	(0.09 to 0.18 lbs ai per Acre)	
Squash	Whiteflies including: Silverleaf/ Sweetpotato (B and Q Biotypes)		
Fruiting Vegetables			
Eggplant			
Peppers			
Tomato			
Head and Stem			
Brassica			
Broccoli			
Brussels Sprouts			
Cabbage			
Cauliflower			
Kohlrabi			

(continued)

**VEGETABLE TRANSPLANTS (IN ENCLOSED STRUCTURES) (continued)
FOLIAR OR BROADCAST SPRAY APPLICATION**

Crops	Pests	Product Rate (By Weight)	Remarks
Leafy Vegetables (Transplants only) (Excluding <i>Brassica</i> spp.)	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including: Silverleaf/ Sweetpotato (B and Q Biotypes)	3.5-5.5 oz per 100 gal 7-11 oz per Acre 0.16-0.25 oz per 1,000 sq ft (0.09 to 0.134 lbs ai per Acre)	Do not make more than one application per crop. Apply only to leafy vegetables being grown as transplants and before transplants are sold. 100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
<p>One (1) level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of <i>Safari</i> 20 SG Insecticide. Begin applications when first pest activity is noticed or when insects reach threshold levels per University/Extension recommendations. Time application before a damaging population becomes established.</p> <p>Restriction: Do not apply more than 1.34 lbs (0.268 lbs ai) per acre of nursery per year. To delay the development of resistance: Do not apply <i>Safari</i> 20 SG Insecticide or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of <i>Safari</i> 20 SG Insecticide or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.</p>			

GREENHOUSE-GROWN TOMATOES*
FOLIAR SPRAY, BROADCAST SPRAY OR SOIL DRENCH APPLICATION

For foliar insect control on tomatoes grown and harvested.

Crop	Pest	Product Rate (By Weight)	Remarks
Tomatoes	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including Silverleaf/ Sweetpotato (B and Q Biotypes)	Foliar Spray 4 to 8 oz (0.05-0.1 lb a.i.) per 100 gal 8-16 oz per Acre (0.1 to 0.2 lbs a.i. per Acre) 0.2-0.4 oz per 1,000 sq ft	100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area. Use Restrictions: Do not use adjuvants or surfactants. Do not apply more than two foliar sprays per crop. Do not apply more than 2.0 lbs (0.4 lb ai) per acre per year as foliar sprays. Do not apply soil drench to plants that have received foliar sprays. Do not apply within one (1) day of harvest.

*Not for use in CA

(continued)

GREENHOUSE-GROWN TOMATOES* (continued)

Crop	Pest	Product Rate (By Weight)	Remarks
Tomatoes	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including Silverleaf/ Sweetpotato (B and Q Biotypes)	Soil Drench 3/4 to 1-1/2 pounds (0.15-0.3 lb a.i.) per 100 gallons 12 to 24 ounces per 100 gallons 1.5-3.0 teaspoons per gallon Apply 4 fl oz of drench solution per gallon of potting media. At 0.75 lb/100 gallons, 360 gallons of drench solution will deliver 2.7 lbs of <i>Safari 20 SG</i> Insecticide per acre. At 1.5 lb/100 gallons, 180 gallons of drench solution will deliver 2.7 lbs of <i>Safari 20 SG</i> Insecticide per acre.	Only apply soil drench to moist soil media. Do not apply to dry or saturated media. Restrictions: Do not leach treated soil media for at least 7 days after application or performance may be reduced. Do not apply more than 2.7 lb (0.54 lb a.i.) per acre per year as a soil drench. Do not use adjuvants or surfactants. Do not apply foliar sprays to plants that have received a soil drench. Do not apply within one (1) day of harvest.

One level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari 20 SG* Insecticide. Begin applications when first pest activity is noticed or when insects reach threshold levels per University/Extension recommendations to maintain control. Time application before a damaging population becomes established.

To delay the development of resistance: Do not apply *Safari 20 SG* Insecticide or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two foliar sprays or one soil drench of *Safari 20 SG* Insecticide or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

*Not for use in CA

ORNAMENTAL PLANTS AND FORESTS – FOLIAR OR BROADCAST SPRAY APPLICATION – OUTDOOR



For foliar insect control on ornamental plants in nurseries, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, reforestation nurseries and forests.

Crops	Pests	Product Rate	Remarks
Ornamental plants including:	Adelgids including: Hemlock Woolly, Balsam Woolly	Foliar Spray	Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days.
Shrubs	Aphids (suppression) including: Balsam, Crepe Myrtle, Green Peach Melon	1/4 to 1/2 lb per 100 gallons	
Bedding Plants	Japanese Beetles (adults)	(4 to 8 oz per 100 gallons)	Tank mixing with a surfactant may improve control of pests such as whitefly, mealybug and scale. Confirm plant safety of tank mix in small area before using on a commercial scale.
Flowering Plants	Lacebugs including: Azalea, Cotoneaster,	(0.05 to 0.1 lbs ai per 100 gallons)	
Foliage Plants	Hawthorne Rhododendron	8-16 oz per Acre	100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Ground Covers	Leaf Beetles, Viburnum	(0.1 to 0.2 lbs ai/A)	
Evergreens	Leafhoppers, including: Glassy-Winged Sharpshooter, Potato	0.2-0.4 oz per 1,000 sq ft	For treatment of small areas:
Ornamental Trees*	Leafminers including: Serpentine	1/2-1.0 tsp per gallon	
Non-Bearing Fruit Trees	Mealybugs including: Citrus, Long-Tailed, Madeira, Obscure, Phormium, Pink Hibiscus		
Non-Bearing Nut Trees	Psyllids including: Asian Citrus		
Non-Bearing Vines	Root Weevils (adults) including: Black Vine, Diaprepes		
Christmas Trees	Sawflies (larvae)		
Trees in Plantations including:	Scale (Armored and Soft) including: Cryptomeria, Cycad Aulacaspis, Elongate Hemlock,		
Conifers	Euonymus, Florida Red, Florida Wax, Tea		
Deciduous trees	Thrips including: Chilli, Gynaikothrips uzeli,		
Reforestation Nurseries	Western Flower (suppression)		
Forests and Wooded Areas: National, Private and State	Whiteflies including: Fig (Ficus), Giant, Greenhouse, Silverleaf / Sweetpotato (B and Q Biotypes)		

ORNAMENTAL PLANTS AND FORESTS – FOLIAR OR BROADCAST SPRAY APPLICATION – OUTDOOR (continued)

One (1) level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari 20 SG Insecticide*. Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days.

Restrictions:

Not for use on house plants grown inside private residences.

Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, landscape or forest per year.

To delay the development of resistance: Do not apply *Safari 20 SG Insecticide* or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of *Safari 20 SG Insecticide* or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

* See species restrictions in the **Application to Ornamental Plants (including Forestry)** section under **Application Information**.

ORNAMENTAL PLANTS – FOLIAR OR BROADCAST SPRAY APPLICATION – INDOOR

For foliar insect control on ornamental plants in greenhouses, interior plantscapes, lath and shadehouses.

Crops	Pests	Product Rate	Remarks
Ornamental plants including:	Adelgids including: Hemlock Woolly, Balsam Woolly	Foliar Spray 1/4 to 1/2 lb	Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days. Tank mixing with a surfactant may improve control of pests such as whitefly, mealybug and scale. Confirm plant safety of tank mix in small area before using on a commercial scale. 100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Shrubs	Aphids (suppression) including: Balsam, Crepe Myrtle, Green Peach Melon	per 100 gallons	
Bedding Plants	Japanese Beetles (adults)	(4 to 8 oz per 100 gallons)	
Flowering Plants	Lacebugs including: Azalea, Cotoneaster, Hawthorne Rhododendron	(0.05 to 0.1 lbs ai per 100 gallons)	
Foliage Plants	Leaf Beetles, Viburnum	8-16 oz per Acre	
Ground Covers	Leafhoppers, including: Glassy-Winged Sharpshooter, Potato	(0.1 to 0.2 lbs ai/A)	
Evergreens	Leafminers including: Serpentine	0.2-0.4 oz per 1,000 sq ft	
Ornamental Trees	Mealybugs including: Citrus, Long-Tailed, Madeira, Obscure, Phormium, Pink Hibiscus	For treatment of small areas:	
Non-Bearing Fruit Trees	Psyllids including: Asian Citrus	1/2-1.0 tsp per gallon	
Non-Bearing Nut Trees	Root Weevils (adults) including: Black Vine, Diaprepes		
Non-Bearing Vines	Sawflies (larvae)		
	Scale (Armored and Soft) including: Cryptomeria, Cycad Aulacaspis, Elongate Hemlock, Euonymus, Florida Red, Florida Wax, Tea		
	Thrips including: Chilli, <i>Gynaikothrips uzeli</i> , Western Flower (suppression)		
	Whiteflies including: Fig (Ficus), Giant, Greenhouse, Silverleaf / Sweetpotato (B and Q Biotypes)		

(continued)

ORNAMENTAL PLANTS – FOLIAR OR BROADCAST SPRAY APPLICATION – INDOOR (continued)

One (1) level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari 20 SG* Insecticide. Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days.

Restrictions:

Not for use on house plants grown inside private residences.

Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery or landscape per year.

To delay the development of resistance: Do not apply *Safari 20 SG* Insecticide or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of *Safari 20 SG* Insecticide or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL: For systemic insect control on containerized and field grown (in-ground) ornamental plants in nurseries, greenhouses, interior plantscapes, lath and shadehouses, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, reforestation nurseries and forests when applied via soil drench, soil injection, micro-irrigation (spaghetti tube or emitter), drip irrigation, overhead irrigation, ebb and flood irrigation equipment or motorized irrigation equipment.

Crops	Pests	Product Rate (By Weight)		Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	Adelgids including: Hemlock Woolly Balsam Woolly	Containerized Plants Soil Media Drench 3/4 to 1-1/2 pounds per 100 gallons 12 to 24 ounces per 100 gallons 1.5-3.0 teaspoons per gallon		Only apply to moist soil media. Do not apply to dry or saturated media. Do not apply media drench until roots from transplanted plugs or liners have extended at least half way to the edge of pots. Do not leach treated soil media for at least 7 days after application or performance may be reduced. Heavy rainfall or excessive irrigation following application may decrease performance. Higher rates will be needed to control insects on woody plants than on herbaceous plants. Poinsettia: For optimal control of whiteflies, treat plants 1-3 weeks after pinch. Late season drenches will take longer to provide effective control.
	Aphids including: Balsam, Crepe Myrtle, Green Peach, Melon	Media Drench Volume for Individual Pots		
		Pot diameter (inches)	Fl oz of dilute solution per pot	
		4	2	
		5	3	
		6	4	
		7	5	
		8	6	
		For larger pot volumes, apply 3-4 fl oz of dilute solution (0.11 to 0.22 g product per 4 fl oz water) per gallon of potting media. Use a drench volume that is sufficient to wet soil media without resulting in overflow or runoff through drain holes in pot.		
		Containerized Plants Media Drench Volume for Plants in Raised Beds, Benches, Bedding Flats, Plug and Liner Trays: Apply sufficient dilute solution to wet soil media without loss of liquid from bottom of bed or liner.		

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)		Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued) Japanese Beetle (adults) Lacebugs including: Azalea Cotoneaster Hawthorne Rhododendron Leaf Beetles including: Elm Viburnum Leafhoppers including: Glass-Winged Sharpshooter Potato Leafminers including: Birch Boxwood Chrysanthemum Holly Serpentine (continued)	Containerized Plants Ebb and Flood Irrigation		Bring several pots to field capacity, let soil dry and then measure amount of water required to bring pots back to field capacity. Multiply the average volume of water required to rehydrate one pot by the number of pots to be treated. Add this volume of water to the minimum amount of water needed to flood the area to be treated. Re-use any returned volume in subsequent irrigation of same plants. For pot diameter greater than 8", use 3.7-7.5 ounces of <i>Safari 20</i> SG Insecticide per 1,000 gallons of potting soil media. Use typical injection ratio for injectors (e.g., 1:100, which equals 1 part injector tank solution: 100 parts irrigation water). Do not mix more than 24 oz of <i>Safari 20</i> SG Insecticide per gallon of injector tank water, or some product may settle out of solution. Calibrate irrigation system to deliver 3-4 fl oz of dilute solution per gallon of potting media.
		Pot diameter (inches)	Ounces per 1,000 pots	
		4	1.9-3.7	
		5	2.8-5.6	
		6	3.7-7.5	
		7	4.7-9.3	
		8	5.6-11.2	
		Chemigation of individual containers using a micro-irrigation system (spaghetti tube)		
		Injection ratio	Ounces per gallon of injector tank water	
		1:100	12-24	

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued) Mealybugs: Citrus Longtailed Madeira Obscure Phormium Pink Hibiscus Root Mimosa Webworm (larvae) Peachtree Borer Pine Tip Moth (larvae) Plantbugs Psyllids including: Asian Citrus Boxwood Root Weevils (larvae and adults) including: Black Vine Diaprepes (continued)	Field Grown (In-Ground) Shrubs 3-6 grams (1.25-2.5 level teaspoons) per foot of height 1.0-2.1 ounces per 10 feet of height	When applied to the soil, <i>Safari 20</i> SG Insecticide is taken up by actively growing trees and shrubs. Speed of control will be dependent on plant size, plant health, environmental conditions and how actively pests are feeding. In actively growing plants, control may be evident within 1-3 weeks after application depending on plant size. Time applications to coincide with when most vulnerable pest life stage is present on plants. Control may be less effective when applied to dry, saturated, or frozen soil, or at times when plants are not actively taking up water from soil. If possible, irrigate dry soils 1-3 days before application, or apply irrigation within 3 days after application. Heavy rainfall or inadequate irrigation immediately following application may decrease performance. Use higher labeled rates for broadleaf evergreens with dense foliage (ex. hollies), and with very large trees. Soil Drench: Mix required dose in water and uniformly apply to soil around base of shrub or tree. Pull back mulch before drenching. Apply 1-4 pints of drench solution per foot of height (shrubs) or inch of trunk diameter (trees). Adjust drench volume based on soil type, soil moisture and thickness of mulch so that product is moved into root zone. To enhance soil penetration in heavy soils and sloping terrain, dig shallow holes around tree or shrub, and apply drench solution in holes. Lower drench volumes may be less effective in dry soils or when applied over heavy mulch unless there is adequate rainfall or irrigation after application to move product into root zone.

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued) Roundheaded Borers (excluding Asian Longhorned) Eucalyptus Longhorned Linden Locust Royal Palm Bug Sawfly larvae Scales (Armored and Soft) including: Azalea Bark Brown Soft Calico California Red Cottony Cushion Cottony Maple Cryptomeria Cycad Aulacaspis Duplachionaspis Elongate Hemlock Euonymus False Florida Red False Oleander (continued)	Field Grown (In-Ground) Trees (Less than 24" diameter at breast height) 3-12 grams (1.25-5.0 level teaspoons) per inch of trunk diameter at breast height (DBH) 1.05-4.2 ounces per 10 inches of trunk diameter at breast height (DBH) For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height. Field Grown (In-Ground) Trees (24" diameter or greater at breast height) 6-12 grams (2.5-5.0 level teaspoons) per inch of trunk diameter at breast height (DBH) 2.1-4.2 ounces per 10 inches of trunk diameter at breast height (DBH) For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height. Hedges 0.25-1.0 oz per foot of hedge height per 100 linear feet of hedge row.	Soil Injection: Mix required dose in water and make at least four injections per shrub or tree with a low-pressure applicator. Use same amount of solution per hole. Injections can be made using the following methods: Grid System – Space injections on a 2.5 ft center extending to drip line. Circle System – Make injections in concentric circles extending inward from drip line. Basal System – Space injections evenly around trunk no more than 24" out from the base. <i>Safari 20</i> SG Insecticide may be soil injected with low volume (e.g. Kioritz injector) or high volume injection equipment. Inject 1-32 fl oz of dilute solution per foot of height or inch of trunk diameter depending on application equipment. Make shallow injections where feeder roots are most concentrated. Apply in enough water to wet the lower 12" of trunk and surrounding soil surface. Apply in a one foot wide band over base of trunk and soil down center of hedgerow. To improve performance, rake back mulch before application.

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)		Remarks
		Row spacing in feet	Ounces per 1,000 linear feet of row	
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	Scales (Armored and Soft) (cont.) Fig (Ficus) Wax Fletcher Florida Red Florida Wax Indian Wax Lecanium Lobate Lac <i>Melanaspis deklei</i> Obscure Oystershell Poplar (Aspen) Pine Needle Tea Tuliptree Spittlebugs Tent Caterpillar (larvae) Thrips including: Chilli (suppression) Citrus Cuban Laurel Gladiolus <i>Gynaikothrips uzeli</i> (suppression) Western Flower (suppression) (continued)	Field Grown Nursery Stock		Apply as a uniform band in row over root zone and lower 6-12" of trunk. Apply from peak adult flight to peak egg hatch. Apply in at least two gallons of water per 1,000 linear feet. Irrigate after application to move product into soil profile. Control any weeds in treated area prior to application, or performance may be reduced. Adjust rates accordingly for other row spacing. Irrigate after application to move <i>Safari</i> 20 SG Insecticide to the root zone. Apply over the top of ornamental plant beds in a water volume sufficient to move product to soil surface. If necessary, irrigate after application to move product off of foliage and into upper root zone of soil. May be less effective on large woody shrubs than on herbaceous annuals and perennials.
		Banded spray application to soil surface (2.7 lbs per acre).		
		3	3	
		4	4	
		5	5	
		6	6	
		7	7	
		8	8	
	Broadcast spray to soil of plant beds (2.7 lbs per acre).			

(continued)

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued) Treehoppers Walnut Twig Beetle Whiteflies including: Ficus Giant Greenhouse Silverleaf / Sweetpotato (B and Q Biotypes) White Grubs including: Oriental Beetle White Pine Weevil		

(continued)

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Important Notes:

One (1) level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari 20 SG* Insecticide.

For all soil applications, including chemigation, retreatments may be made after 7 days but do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, landscape or forest per year.

Restrictions:

Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, landscape or forest per year.

To delay the development of resistance in greenhouses, nurseries and interiorscapes, do not make more than one soil application per crop cycle or three months, whichever is shorter. Refer to "Resistance Management" section of the label for additional guidelines.

*See species restrictions in the **Application to Ornamental Plants (including Forestry)** section under **Application Information**.

ORNAMENTAL PLANTS AND FORESTS

BASAL TRUNK SPRAYS IN TREES AND LARGE SHRUBS

For systemic insect control in containerized and field grown (in-ground) ornamental trees and shrubs in nurseries, interior landscapes, lath and shadehouses, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations and forests when applied as a trunk spray.

Crops	Pests	Product Rate (By Weight)	Remarks
Shrubs Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Trees in Plantations including: Conifer Deciduous Reforestation Nurseries Forests and Wooded Areas: National, Private and State	Adelgids including: Hemlock Woolly Aphids Flatheaded Borers including: Alder Bronze Birch Emerald Ash Flatheaded Appletree Two-lined Chestnut Lacebugs Leaf Beetles Leafhoppers Leafminers Mealybugs Mountain Pine Beetle (continued)	12-24 oz per gallon Depending on bark type and thickness, one gallon of spray solution will typically cover 65-85" of cumulative trunk diameter (1.5-2.0 fl oz per inch of trunk diameter) when applied to trunk between soil surface and 4-5 feet above soil surface.	When sprayed on the trunk, <i>Safari 20 SG</i> Insecticide will be absorbed through the bark and into the vascular system, and then transported throughout the tree. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively transpiring trees, control may be evident within 1-3 weeks after application. Spray bark on root flare (buttress roots) and on trunk between soil surface and 4-5 feet above the soil surface. Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off onto soil. Apply ONLY with a low volume sprayer operated at less than 20 PSI to prevent tree damage, bounce back and drift of spray droplets. Time applications to coincide with when most vulnerable pest life stage is present on plants. Do not apply to wet bark, during rainfall or if rain is expected within 12 hours. Control may be less effective in trees with thick bark, and at times when trees are not actively growing or transpiring. For Mountain Pine Beetle: apply from 2 weeks before to 2 weeks after expected peak of adult flight activity.

ORNAMENTAL PLANTS AND FORESTS (continued)
BASAL TRUNK SPRAYS IN TREES AND LARGE SHRUBS (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Shrubs Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Trees in Plantations including: Conifer Deciduous Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued) Pine Tip Moth (larvae) Psyllids Roundheaded Borers (excluding Asian Longhorned) Scales including: Calico Cryptomeria Elongate Hemlock Fig (Ficus) Wax Thrips (suppression) Walnut Twig Beetle Whiteflies including: Fig (Ficus)	12-24 oz per gallon Depending on bark type and thickness, one gallon of spray solution will typically cover 65-85" of cumulative trunk diameter (1.5-2.0 fl oz per inch of trunk diameter) when applied to trunk between soil surface and 4-5 feet above soil surface.	When sprayed on the trunk, <i>Safari</i> 20 SG Insecticide will be absorbed through the bark and into the vascular system, and then transported throughout the tree. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively transpiring trees, control may be evident within 1-3 weeks after application. Spray bark on root flare (buttress roots) and on trunk between soil surface and 4-5 feet above the soil surface. Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off onto soil. Apply ONLY with a low volume sprayer operated at less than 20 PSI to prevent tree damage, bounce back and drift of spray droplets. Time applications to coincide with when most vulnerable pest life stage is present on plants. Do not apply to wet bark, during rainfall or if rain is expected within 12 hours. Control may be less effective in trees with thick bark, and at times when trees are not actively growing or transpiring. For Mountain Pine Beetle: apply from 2 weeks before to 2 weeks after expected peak of adult flight activity.

(continued)

ORNAMENTAL PLANTS AND FORESTS (continued)
BASAL TRUNK SPRAYS IN TREES AND LARGE SHRUBS (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Christmas Trees Ornamental Trees* with trunk diameter less than 3" at soil line	Elongate Hemlock Scale Cryptomeria Scale Ficus (Fig) Whitefly	1.5-6.0 oz/gallon One gallon of spray solution will typically cover 325-425" of cumulative trunk diameter (0.3-0.4 fl oz per inch of trunk diameter) when applied to trunk between soil surface and 1 foot above soil surface.	For Christmas trees and ornamental trees less than 3" in diameter at soil line, spray trunk just to point of runoff between soil surface and 12" above soil surface.
<p>One (1) level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz by weight of <i>Safari</i> 20 SG Insecticide. Restrictions: Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, forest or landscape per year. * See species restrictions in the Application to Ornamental Plants (including Forestry) section under Application Information.</p>			

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool dry place. Do not store diluted spray. For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-CLEANUP.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container: Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration.

CONDITIONS OF SALE

Valent U.S.A. LLC warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions to the crops specified. To the extent consistent with applicable law, there are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

©2019 Valent U.S.A. LLC

Safari is a registered trademark of Valent U.S.A. LLC

Manufactured for

Valent U.S.A. LLC

P.O. Box 8025

Walnut Creek CA 94596-8025

Made in U.S.A.

Form 1510-I

TAJ 15AUG16


EPA Reg. No. 86203-11-59639

EPA Est. 67545-AZ-1

Safari®

20 SG INSECTICIDE

GROUP 4A INSECTICIDE



FOR FOLIAR AND SYSTEMIC INSECT CONTROL IN ORNAMENTAL PLANTS AND VEGETABLE TRANSPLANTS IN ENCLOSED STRUCTURES.

FOR GREENHOUSE, NURSERY, INTERIOR PLANTSCAPE, OUTDOOR LANDSCAPE, GREENHOUSE TOMATO AND FORESTRY USE ONLY.

Active Ingredient:
Dinotefuran, [N-methyl-N'-nitro-N''-(
(tetrahydro-3-furanyl)methyl)guanidine]. . . 20%
Other Ingredients. 80%
Total 100%
EPA Reg. No. 86203-11-59639
EPA Est. 67545-AZ-1

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

**SEE INSIDE BOOKLET FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**


NET CONTENTS 3 POUNDS

Form 1510-1

GROUP 4A INSECTICIDE

Safari®

20 SG INSECTICIDE



FOR FOLIAR AND SYSTEMIC INSECT CONTROL IN ORNAMENTAL PLANTS AND VEGETABLE TRANSPLANTS IN ENCLOSED STRUCTURES.

FOR GREENHOUSE, NURSERY, INTERIOR PLANTSCAPE, OUTDOOR LANDSCAPE AND FORESTRY USE ONLY.

Active Ingredient:
 Dinotefuran, [N-methyl-N'-nitro-N''-
 ((tetrahydro-3-furanyl)methyl)guanidine].. 20%
 Other Ingredients 80%
 Total 100%

EPA Reg. No. 86203-11-59639
 EPA Est. 67545-AZ-1

NET CONTENTS 12 OUNCES

KEEP OUT OF REACH OF CHILDREN CAUTION
SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS



Form 1812-F

FIRST AID

- If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
- If swallowed:** Call poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by the poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
- If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
- If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Causes moderate eye irritation. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made out of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or viton \geq 14 mils

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not dispose equipment washwaters or rinsate into a natural drain or water body.

This product is toxic to honey bees. The persistence of residues and potential residual toxicity of dinotefuran in nectar and pollen suggests the possibility of chronic toxic risk to honey bee larvae and the eventual instability of the hive.

- This product is toxic to bees exposed to residues for more than 38 hours following treatment.
- Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

Dinotefuran and its degradate, MNG, have the properties and characteristics associated with chemicals detected in groundwater. The high water solubility of dinotefuran, and its degradate, MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

SPRAY DRIFT ADVISORY

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered for sale, use or consumption.

PROTECTION OF POLLINATORS APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site pollinator attractive habitat can result in bee kills.

(continued)

PROTECTION OF POLLINATORS (continued)

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>. Pesticide incidents (for example, bee kills) should immediately be reported to the State/Tribal lead agency. For contact information for your State, go to: www.aapco.org. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

FOR COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS



- Do not apply this product while bees are foraging.
- This product is toxic to bees exposed to residue for more than 38 hours following treatment.
- Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

NON-AGRICULTURAL USES



Do not apply Safari® 20 SG Insecticide while bees are foraging. Do not apply Safari 20 SG Insecticide to plants that are flowering. Only apply after all flower petals have fallen off.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural insecticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

EXCEPTION: If product is drenched or soil-injected, workers may enter the area at any time if there will be no contact with anything that has been treated.

(continued)

(continued)

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves (made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or viton \geq 14 mils)

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow others to enter treated areas until sprays have dried.

APPLICATION INFORMATION

- Applications of *Safari 20 SG* Insecticide in residential areas may be made by commercially licensed applicators.

Application to Ornamental Plants (including Forestry):

- *Safari 20 SG* Insecticide can be applied as a foliar spray, a broadcast spray, a soil drench, soil injection and via chemigation for insect control in ornamental plants in greenhouses, nurseries, outdoor landscapes and interior plantscapes.
- *Safari 20 SG* Insecticide is a systemic product and will be taken up by the root system and translocated upward throughout the plant. When applied as a foliar spray, the product offers translaminar and locally systemic control of foliar pests.

- When applied to the soil, *Safari 20 SG* Insecticide will be translocated more quickly in herbaceous plants than in woody shrubs and trees. Speed of insect control will range from as little as one day for small herbaceous plants in containers, to several weeks in large trees growing in the landscape.
- Do not apply more than a total of 2.7 lbs of product (0.54 lb active ingredient) per acre per year for all application types.
- **Do not apply this product, by any application method, to linden, basswood or other *Tilia* species.**
- **Use on *Myoporum laetum* is not recommended.**

Application to Vegetable Transplants:

- *Safari 20 SG* Insecticide can be applied as a foliar spray or a broadcast spray for insect control in vegetable transplants.
- Do not apply more than 1.34 lbs (0.268 lbs ai) per acre of nursery per year.

MIXING INSTRUCTIONS:

Safari 20 SG Insecticide Alone: Add half of the required amount of water to the mix tank. With the agitator running, add the desired amount of *Safari 20 SG* Insecticide to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after *Safari 20 SG* Insecticide has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Safari 20 SG Insecticide + Tank Mixtures: Add half of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, emulsifiable concentrates, and surfactants/adjuvants. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

NOTE: When using *Safari* 20 SG Insecticide in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including *Safari* 20 SG Insecticide. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using *Safari* 20 SG Insecticide in a tank mixture, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Do not exceed label dosage rate. Do not mix this product with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Volumes, weights, and conversions

Use the following equivalent, unpacked volumetric weights

1 tablespoon = 0.22 oz = approximately 1/4 oz;

2.5 tablespoons = 0.55 oz = approximately 1/2 oz;

3.5 tablespoons = 0.77 oz = approximately 3/4 oz;

4.5 tablespoons = 0.99 oz = approximately 1 oz;

28 grams = 1.0 oz

2.7 lb = 43.2 oz

Compatibility

IMPORTANT: The safety of all potential tank mixes has not been tested on all crops. Before applying any tank mixture not specifically listed on this label, confirm the safety to the target crop.

Safari 20 SG Insecticide is compatible with most commonly used pesticides, crop oils, adjuvants, and nutritional sprays. However, since it is not possible to test all possible

mixtures, pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with *Safari* 20 SG Insecticide. To determine the physical compatibility of *Safari* 20 SG Insecticide with other products, use a jar test, as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for additional required ingredients to the spray tank.

RESISTANCE MANAGEMENT

Safari 20 SG Insecticide contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly in the same crop or in successive years as the primary method of control for a targeted species. This may result in partial or total loss of control of those species by *Safari* 20 SG Insecticide or other Group 4A insecticides.

To delay the development of insecticide resistance in greenhouse, nursery and interiorscape use sites, strongly consider the following guidelines:

- Do not apply *Safari* 20 SG Insecticide or other Group 4A insecticides to consecutive generations of the same insect pest species.
- Do not drench soil media with *Safari* 20 SG Insecticide or other Group 4A insecticides more than one time per crop cycle or three months, whichever is shorter.
- Do not make more than two foliar or broadcast sprays of *Safari* 20 SG Insecticide or other Group 4A insecticides to a single crop during a two-month period.

(continued)

(continued)

- Do not make more than one soil drench and one foliar or broadcast spray with *Safari* 20 SG Insecticide or other Group 4A insecticides during a two-month period.
- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturers for insecticide resistance management and/or IPM guidelines for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Valent U.S.A. LLC, at toll free number: 1-800-898-2536.

APPLICATION PROCEDURES AND SPRAY EQUIPMENT

Ground Application: Select spray nozzles that will provide accurate and uniform spray deposition. Use spray nozzles that provide medium-sized droplets and reduce drift. To help insure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult nozzle manufacturers and/or State Extension Service specialists.

Apply *Safari* 20 SG Insecticide using sufficient water volume to provide thorough and uniform coverage. In situations where a dense canopy exists and/or pest pressure is high, use greater water volumes. The use of a spray adjuvant may improve spray coverage. Do not apply under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Applications to ornamental plants, forestry and vegetable transplants: *Safari* 20 SG Insecticide can be applied using many different types of application equipment. Apply in sufficient water to ensure good coverage of ornamental plants. Tank mixing with a surfactant will produce better coverage when making applications to plants with hard to wet foliage such as holly or pine. If concentrate or mist type spray equipment

is used, apply the same amount of product on the sprayed area as would be used in a dilute solution. To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. Applications can be made to foliage or as a soil drench.

RESTRICTIONS

- With the exception of non-livestock animals, do not graze treated areas or use clippings from treated areas for feed or forage.
- Prevent runoff or puddling of irrigation water following application.
- Keep children and pets off treated areas until spray has dried.
- Do not apply to areas that are water logged or saturated, or frozen, which will not allow penetration into the root zone of the plant.

APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION):

Safari 20 SG Insecticide may be applied by injection into an irrigation system, either alone or in combination with other pesticides or chemicals that are registered for application through irrigation systems. Dilution ratios are normally 1:100 to 1:200, depending on the system. Apply this product only through microirrigation (individual spaghetti tube), drip irrigation, overhead irrigation, or motorized calibrated irrigation equipment (Ornamentals). Do not apply through any other type of irrigation system. Lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make adjustments when necessary.

Using Water from Public Water Systems:

DO NOT APPLY *SAFARI* 20 SG INSECTICIDE THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year. *Safari 20 SG* Insecticide may be applied through irrigation systems which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements:

Operating Instructions for Irrigation Systems:

1. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
2. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

6. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Do not apply when wind speed favors drift beyond the area intended.

Calibration and Application Instructions:

Apply *Safari 20 SG* Insecticide under the schedule specified in the specific use instructions, not according to the irrigation schedule unless the events coincide. In general, set the equipment to apply the minimum amount of water per acre. Run the system at 86-90% of the manufacturer's maximum rated travel speed.

The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Check with State and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

MINIMIZING SPRAY DRIFT

As with all crop protection products, it is important to minimize off-target movement. Do not allow spray to drift onto adjacent land, crops, or aquatic areas. To minimize spray drift:

1. Make applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 10 mph. Do not apply when wind gusts approach 10 mph.
2. Risk of exposure to sensitive aquatic areas can be reduced by not applying when wind direction is toward the aquatic area.
3. Do not cultivate or plant crops within 25 feet of the aquatic area as to allow growth of a vegetative filter strip.

4. Do not make applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with increased height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
5. Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by not using excessive spray boom pressure.
6. Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage. Do not apply more than 10 ft above the crop canopy.
7. For aerial applications, mount spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use minimum practical boom length and do not use boom that exceeds 75% of wing span or rotor diameter.

Air Assisted (Air Blast) Tree and Vine Sprayers (Ornamentals Only):

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift:

1. Adjust deflectors and aiming devices so that spray is only directed into the canopy.
2. Block off upward pointed nozzles when there is no overhanging canopy.
3. Use only enough air volume to penetrate the canopy and provide good coverage. Use a minimum of 50 gallons finished spray per acre.
4. Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.

**VEGETABLE TRANSPLANTS (IN ENCLOSED STRUCTURES)
FOLIAR OR BROADCAST SPRAY APPLICATION**

For foliar insect control on vegetable transplants grown in enclosed structures.

Crops	Pests	Product Rate (By Weight)	Remarks
Cucurbits (Transplants only)	Aphids	3.5-7.0 oz per 100 gal	Do not make more than one application per crop.
Cantaloupe	Leafminers	7-14 oz per Acre	Apply only to cucurbits and brassica being grown as transplants and before transplants are sold.
Cucumber	Mealybugs	0.16-0.32 oz per 1,000 sq ft	100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Melons	Thrips (suppression)	(0.09 to 0.18 lbs ai per Acre)	
Squash	Whiteflies including:		
Fruiting Vegetables	Silverleaf/		
Eggplant	Sweetpotato		
Peppers	(B and Q Biotypes)		
Tomato			
Head and Stem			
Brassica			
Broccoli			
Brussels Sprouts			
Cabbage			
Cauliflower			
Kohlrabi			

(continued)

VEGETABLE TRANSPLANTS (IN ENCLOSED STRUCTURES) (continued)
FOLIAR OR BROADCAST SPRAY APPLICATION

Crops	Pests	Product Rate (By Weight)	Remarks
Leafy Vegetables (Transplants only) (Excluding <i>Brassica</i> spp.)	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including: Silverleaf/ Sweetpotato (B and Q Biotypes)	3.5-5.5 oz per 100 gal 7-11 oz per Acre 0.16-0.25 oz per 1,000 sq ft (0.09 to 0.134 lbs ai per Acre)	Do not make more than one application per crop. Apply only to leafy vegetables being grown as transplants and before transplants are sold. 100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.

One (1) level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari* 20 SG Insecticide.

Begin applications when first pest activity is noticed or when insects reach threshold levels per University/Extension recommendations. Time application before a damaging population becomes established.

Restriction:

Do not apply more than 1.34 lbs (0.268 lbs ai) per acre of nursery per year.

To delay the development of resistance: Do not apply *Safari* 20 SG Insecticide or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of *Safari* 20 SG Insecticide or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

GREENHOUSE-GROWN TOMATOES*
FOLIAR SPRAY, BROADCAST SPRAY OR SOIL DRENCH APPLICATION

For foliar insect control on tomatoes grown and harvested.

Crop	Pest	Product Rate (By Weight)	Remarks
Tomatoes	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including: Silverleaf/ Sweetpotato (B and Q Biotypes)	Foliar Spray 4 to 8 oz (0.05-0.1 lb a.i.) per 100 gal 8-16 oz per Acre (0.1 to 0.2 lbs a.i. per Acre) 0.2-0.4 oz per 1,000 sq ft	100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area. Use Restrictions: Do not use adjuvants or surfactants. Do not apply more than two foliar sprays per crop. Do not apply more than 2.0 lbs (0.4 lb ai) per acre per year as foliar sprays. Do not apply soil drench to plants that have received foliar sprays. Do not apply within one (1) day of harvest.

*Not for use in CA

(continued)

GREENHOUSE-GROWN TOMATOES* (continued)

Crop	Pest	Product Rate (By Weight)	Remarks
Tomatoes	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including: Silverleaf/Sweetpotato (B and Q Biotypes)	Soil Drench 3/4 to 1-1/2 pounds (0.15-0.3 lb a.i.) per 100 gallons 12 to 24 ounces per 100 gallons 1.5-3.0 teaspoons per gallon Apply 4 fl oz of drench solution per gallon of potting media. At 0.75 lb/100 gallons, 360 gallons of drench solution will deliver 2.7 lbs of <i>Safari 20 SG</i> Insecticide per acre. At 1.5 lb/100 gallons, 180 gallons of drench solution will deliver 2.7 lbs of <i>Safari 20 SG</i> Insecticide per acre.	Only apply soil drench to moist soil media. Do not apply to dry or saturated media. Restrictions: Do not leach treated soil media for at least 7 days after application or performance may be reduced. Do not apply more than 2.7 lb (0.54 lb a.i.) per acre per year as a soil drench. Do not use adjuvants or surfactants. Do not apply foliar sprays to plants that have received a soil drench. Do not apply within one (1) day of harvest.

One level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari 20 SG* Insecticide.

Begin applications when first pest activity is noticed or when insects reach threshold levels per University/Extension recommendations to maintain control. Time application before a damaging population becomes established.

To delay the development of resistance: Do not apply *Safari 20 SG* Insecticide or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two foliar sprays or one soil drench of *Safari 20 SG* Insecticide or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

*Not for use in CA

ORNAMENTAL PLANTS AND FORESTS – FOLIAR OR BROADCAST SPRAY APPLICATION – OUTDOOR



For foliar insect control on ornamental plants in nurseries, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, reforestation nurseries and forests.

Crops	Pests	Product Rate	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	Adelgids including: Hemlock Woolly, Balsam Woolly Aphids (suppression) including: Balsam, Crepe Myrtle, Green Peach Melon Japanese Beetles (adults) Lacebugs including: Azalea, Cotoneaster, Hawthorne Rhododendron Leaf Beetles, Viburnum Leafhoppers, including: Glassy-Winged Sharpshooter, Potato Leafminers including: Serpentine Mealybugs including: Citrus, Long-Tailed, Madeira, Obscure, Phormium, Pink Hibiscus Psyllids including: Asian Citrus Root Weevils (adults) including: Black Vine, Diaprepes Sawflies (larvae) Scale (Armored and Soft) including: Cryptomeria, Cycad Aulacaspis, Elongate Hemlock, Euonymus, Florida Red, Florida Wax, Tea Thrips including: Chilli, Gynaikothrips uzeli, Western Flower (suppression) Whiteflies including: Fig (Ficus), Giant, Greenhouse, Silverleaf / Sweetpotato (B and Q Biotypes)	Foliar Spray 1/4 to 1/2 lb per 100 gallons (4 to 8 oz per 100 gallons) (0.05 to 0.1 lbs ai per 100 gallons) 8-16 oz per Acre (0.1 to 0.2 lbs ai/A) 0.2-0.4 oz per 1,000 sq ft For treatment of small areas: 1/2-1.0 tsp per gallon	Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days. Tank mixing with a surfactant may improve control of pests such as whitefly, mealybug and scale. Confirm plant safety of tank mix in small area before using on a commercial scale. 100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.

ORNAMENTAL PLANTS AND FORESTS – FOLIAR OR BROADCAST SPRAY APPLICATION – OUTDOOR (continued)

One (1) level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari* 20 SG Insecticide.

Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days.

Restrictions:

Not for use on house plants grown inside private residences.

Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, landscape or forest per year.

To delay the development of resistance: Do not apply *Safari* 20 SG Insecticide or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of *Safari* 20 SG Insecticide or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

*See species restrictions in the **Application to Ornamental Plants (including Forestry)** section under **Application Information**.

ORNAMENTAL PLANTS – FOLIAR OR BROADCAST SPRAY APPLICATION – INDOOR

For foliar insect control on ornamental plants in greenhouses, interior plantscapes, lath and shadehouses.

Crops	Pests	Product Rate	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines	Adelgids including: Hemlock Woolly, Balsam Woolly Aphids (suppression) including: Balsam, Crepe Myrtle, Green Peach Melon Japanese Beetles (adults) Lacebugs including: Azalea, Cotoneaster, Hawthorne, Rhododendron Leaf Beetles, Viburnum Leafhoppers, including: Glassy-Winged Sharpshooter, Potato Leafminers including: Serpentine Mealybugs including: Citrus, Long-Tailed, Madeira, Obscure, Phormium, Pink Hibiscus Psyllids including: Asian Citrus Root Weevils (adults) including: Black Vine, Diaprepes Sawflies (larvae) Scale (Armored and Soft) including: Cryptomeria, Cycad Aulacaspis, Elongate Hemlock, Euonymus, Florida Red, Florida Wax, Tea Thrips including: Chilli, <i>Gynaikothrips uzeli</i> , Western Flower (suppression) Whiteflies including: Fig (Ficus), Giant, Greenhouse, Silverleaf / Sweetpotato (B and Q Biotypes)	Foliar Spray 1/4 to 1/2 lb per 100 gallons (4 to 8 oz per 100 gallons) (0.05 to 0.1 lbs ai per 100 gallons) 8-16 oz per Acre (0.1 to 0.2 lbs ai/A) 0.2-0.4 oz per 1,000 sq ft For treatment of small areas: 1/2-1.0 tsp per gallon	Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days. Tank mixing with a surfactant may improve control of pests such as whitefly, mealybug and scale. Confirm plant safety of tank mix in small area before using on a commercial scale. 100 gals of spray mix will treat 20,000 sq ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.

ORNAMENTAL PLANTS – FOLIAR OR BROADCAST SPRAY APPLICATION – INDOOR (continued)

One (1) level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari* 20 SG Insecticide.

Make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days.

Restrictions:

Not for use on house plants grown inside private residences.

Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery or landscape per year.

To delay the development of resistance: Do not apply *Safari* 20 SG Insecticide or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of *Safari* 20 SG Insecticide or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL: For systemic insect control on containerized and field grown (in-ground) ornamental plants in nurseries, greenhouses, interior plantscapes, lath and shadehouses, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, reforestation nurseries and forests when applied via soil drench, soil injection, micro-irrigation (spaghetti tube or emitter), drip irrigation, overhead irrigation, ebb and flood irrigation equipment or motorized irrigation equipment.

Crops	Pests	Product Rate (By Weight)		Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	Adelgids including: Hemlock Woolly Balsam Woolly Aphids including: Balsam, Crepe Myrtle, Green Peach, Melon Bagworms Eastern Tent Caterpillar Erythrina Gall Wasp Flatheaded Borers including: Alder, Bronze Birch, Emerald Ash, Flatheaded Appletree, Two-Lined Chestnut Froghoppers Fungus Gnats (larvae) Gypsy Moth (larvae) Horned Oak Gall (continued)	Containerized Plants Soil Media Drench 3/4 to 1-1/2 pounds per 100 gallons 12 to 24 ounces per 100 gallons 1.5-3.0 teaspoons per gallon		Only apply to moist soil media. Do not apply to dry or saturated media. Do not apply media drench until roots from transplanted plugs or liners have extended at least half way to the edge of pots. Do not leach treated soil media for at least 7 days after application or performance may be reduced. Heavy rainfall or excessive irrigation following application may decrease performance. Higher rates will be needed to control insects on woody plants than on herbaceous plants. Poinsettia: For optimal control of whiteflies, treat plants 1-3 weeks after pinch. Late season drenches will take longer to provide effective control.
	Media Drench Volume for Individual Pots			
	Pot diameter (inches)	Fl oz of dilute solution per pot		
	4	2		
	5	3		
	6	4		
	7	5		
	8	6		
	For larger pot volumes, apply 3-4 fl oz of dilute solution (0.11 to 0.22 g product per 4 fl oz water) per gallon of potting media. Use a drench volume that is sufficient to wet soil media without resulting in overflow or runoff through drain holes in pot.			
	Containerized Plants Media Drench Volume for Plants in Raised Beds, Benches, Bedding Flats, Plug and Liner Trays: Apply sufficient dilute solution to wet soil media without loss of liquid from bottom of bed or liner.			

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)		Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued)	Containerized Plants Ebb and Flood Irrigation		Bring several pots to field capacity, let soil dry and then measure amount of water required to bring pots back to field capacity. Multiply the average volume of water required to rehydrate one pot by the number of pots to be treated. Add this volume of water to the minimum amount of water needed to flood the area to be treated. Re-use any returned volume in subsequent irrigation of same plants. For pot diameter greater than 8", use 3.7-7.5 ounces of <i>Safari 20 SG</i> Insecticide per 1,000 gallons of potting soil media. Use typical injection ratio for injectors (e.g., 1:100, which equals 1 part injector tank solution: 100 parts irrigation water). Do not mix more than 24 oz of <i>Safari 20 SG</i> Insecticide per gallon of injector tank water, or some product may settle out of solution. Calibrate irrigation system to deliver 3-4 fl oz of dilute solution per gallon of potting media.
	Japanese Beetle (adults)	Pot diameter (inches)	Ounces per 1,000 pots	
	Lacebugs including:	4	1.9-3.7	
	Azalea	5	2.8-5.6	
	Cotoneaster	6	3.7-7.5	
	Hawthorne	7	4.7-9.3	
	Rhododendron	8	5.6-11.2	
	Leaf Beetles including:	Chemigation of individual containers using a micro-irrigation system (spaghetti tube)		
	Elm	Injection ratio	Ounces per gallon of injector tank water	
	Viburnum	1:100	12-24	

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued)	Field Grown (In-Ground) Shrubs 3-6 grams (1.25-2.5 level teaspoons) per foot of height 1.0-2.1 ounces per 10 feet of height	When applied to the soil, <i>Safari 20 SG</i> Insecticide is taken up by actively growing trees and shrubs. Speed of control will be dependent on plant size, plant health, environmental conditions and how actively pests are feeding. In actively growing plants, control may be evident within 1-3 weeks after application depending on plant size. Time applications to coincide with when most vulnerable pest life stage is present on plants. Control may be less effective when applied to dry, saturated, or frozen soil, or at times when plants are not actively taking up water from soil. If possible, irrigate dry soils 1-3 days before application, or apply irrigation within 3 days after application. Heavy rainfall or inadequate irrigation immediately following application may decrease performance. Use higher labeled rates for broadleaf evergreens with dense foliage (ex. hollies), and with very large trees. Soil Drench: Mix required dose in water and uniformly apply to soil around base of shrub or tree. Pull back mulch before drenching. Apply 1-4 pints of drench solution per foot of height (shrubs) or inch of trunk diameter (trees). Adjust drench volume based on soil type, soil moisture and thickness of mulch so that product is moved into root zone. To enhance soil penetration in heavy soils and sloping terrain, dig shallow holes around tree or shrub, and apply drench solution in holes. Lower drench volumes may be less effective in dry soils or when applied over heavy mulch unless there is adequate rainfall or irrigation after application to move product into root zone.
	Mealybugs:		
	Citrus		
	Longtailed		
	Madeira		
	Obscure		
	Phormium		
	Pink Hibiscus		
	Root		
	Mimosa		
	Webworm (larvae)		
	Peachtree		
	Borer		
	Pine Tip Moth (larvae)		
	Plantbugs		
	Psyllids including:		
	Asian Citrus		
	Boxwood		
	Root Weevils (larvae and adults) including:		
	Black Vine		
Diaprepes (continued)			

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued) Roundheaded Borers (excluding Asian Longhorned) Eucalyptus Longhorned Linden Locust Royal Palm Bug Sawfly larvae Scales (Armored and Soft) including: Azalea Bark Brown Soft Calico California Red Cottony Cushion Cottony Maple Cryptomeria Cycad Aulacaspis Duplacionaspis Elongate Hemlock Euonymus False Florida Red False Oleander (continued)	Field Grown (In-Ground) Trees (Less than 24" diameter at breast height) 3-12 grams (1.25-5.0 level teaspoons) per inch of trunk diameter at breast height (DBH) 1.05-4.2 ounces per 10 inches of trunk diameter at breast height (DBH) For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.	Soil Injection: Mix required dose in water and make at least four injections per shrub or tree with a low-pressure applicator. Use same amount of solution per hole. Injections can be made using the following methods: Grid System – Space injections on a 2.5 ft center extending to drip line. Circle System – Make injections in concentric circles extending inward from drip line. Basal System – Space injections evenly around trunk no more than 24" out from the base. Safari 20 SG Insecticide may be soil injected with low volume (e.g. Kioritz injector) or high volume injection equipment. Inject 1-32 fl oz of dilute solution per foot of height or inch of trunk diameter depending on application equipment. Make shallow injections where feeder roots are most concentrated.
		Field Grown (In-Ground) Trees (24" diameter or greater at breast height) 6-12 grams (2.5-5.0 level teaspoons) per inch of trunk diameter at breast height (DBH) 2.1-4.2 ounces per 10 inches of trunk diameter at breast height (DBH) For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.	
		Hedges 0.25-1.0 oz per foot of hedge height per 100 linear feet of hedge row.	

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)	Remarks														
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Christmas Trees Trees in Plantations including: Conifers Deciduous Trees Reforestation Nurseries Forests and Wooded Areas: National, Private and State	Scales (Armored and Soft) (cont.) Fig (Ficus) Wax Fletcher Florida Red Florida Wax Indian Wax Lecanium Lobate Lac <i>Melanaspis deklei</i> Obscure Oystershell Poplar (Aspen) Pine Needle Tea Tuliptree Spittlebugs Tent Caterpillar (larvae) Thrips including: Chilli (suppression) Citrus Cuban Laurel Gladiolus <i>Gynaikothrips uzeli</i> (suppression) Western Flower (suppression) (continued)	Field Grown Nursery Stock Banded spray application to soil surface (2.7 lbs per acre).	Apply as a uniform band in row over root zone and lower 6-12" of trunk. Apply from peak adult flight to peak egg hatch. Apply in at least two gallons of water per 1,000 linear feet. Irrigate after application to move product into soil profile. Control any weeds in treated area prior to application, or performance may be reduced. Adjust rates accordingly for other row spacing. Irrigate after application to move Safari 20 SG Insecticide to the root zone. Apply over the top of ornamental plant beds in a water volume sufficient to move product to soil surface. If necessary, irrigate after application to move product off of foliage and into upper root zone of soil. May be less effective on large woody shrubs than on herbaceous annuals and perennials.														
		<table border="1"> <tr> <th>Row spacing in feet</th> <th>Ounces per 1,000 linear feet of row</th> </tr> <tr> <td>3</td> <td>3</td> </tr> <tr> <td>4</td> <td>4</td> </tr> <tr> <td>5</td> <td>5</td> </tr> <tr> <td>6</td> <td>6</td> </tr> <tr> <td>7</td> <td>7</td> </tr> <tr> <td>8</td> <td>8</td> </tr> </table>		Row spacing in feet	Ounces per 1,000 linear feet of row	3	3	4	4	5	5	6	6	7	7	8	8
		Row spacing in feet		Ounces per 1,000 linear feet of row													
		3		3													
		4		4													
		5		5													
		6		6													
		7		7													
		8		8													
		Broadcast spray to soil of plant beds (2.7 lbs per acre).															

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Ornamental plants including:	(continued)		
Shrubs	Treehoppers		
Bedding Plants	Walnut Twig Beetle		
Flowering Plants	Whiteflies including:		
Foliage Plants	Ficus		
Ground Covers	Giant		
Evergreens	Greenhouse		
Ornamental Trees*	Silverleaf /		
Non-Bearing	Sweetpotato		
Fruit Trees	(B and Q		
Non-Bearing	Biotypes)		
Nut Trees	White Grubs		
Non-Bearing Vines	including:		
Christmas Trees	Oriental Beetle		
Trees in Plantations including:	White Pine Weevil		
Conifers			
Deciduous Trees			
Reforestation Nurseries			
Forests and Wooded Areas: National, Private and State			

(continued)

ORNAMENTAL PLANTS AND FORESTS – APPLICATION TO SOIL (continued)

Important Notes:

One (1) level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari* 20 SG Insecticide.

For all soil applications, including chemigation, retreatments may be made after 7 days but do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, landscape or forest per year.

Restrictions:

Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, landscape or forest per year.

To delay the development of resistance in greenhouses, nurseries and interiorscapes, do not make more than one soil application per crop cycle or three months, whichever is shorter. Refer to "Resistance Management" section of the label for additional guidelines.

*See species restrictions in the **Application to Ornamental Plants (including Forestry)** section under **Application Information**.

ORNAMENTAL PLANTS AND FORESTS

BASAL TRUNK SPRAYS IN TREES AND LARGE SHRUBS

For systemic insect control in containerized and field grown (in-ground) ornamental trees and shrubs in nurseries, interior plantscapes, lath and shadehouses, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations and forests when applied as a trunk spray.

Crops	Pests	Product Rate (By Weight)	Remarks
Shrubs Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Trees in Plantations including: Conifer Deciduous Reforestation Nurseries Forests and Wooded Areas: National, Private and State	Adelgids including: Hemlock Woolly Aphids Flatheaded Borers including: Alder Bronze Birch Emerald Ash Flatheaded Appletree Two-lined Chestnut Lacebugs Leaf Beetles Leafhoppers Leafminers Mealybugs Mountain Pine Beetle (continued)	12-24 oz per gallon Depending on bark type and thickness, one gallon of spray solution will typically cover 65-85" of cumulative trunk diameter (1.5-2.0 fl oz per inch of trunk diameter) when applied to trunk between soil surface and 4-5 feet above soil surface.	When sprayed on the trunk, <i>Safari 20 SG</i> Insecticide will be absorbed through the bark and into the vascular system, and then transported throughout the tree. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively transpiring trees, control may be evident within 1-3 weeks after application. Spray bark on root flare (buttress roots) and on trunk between soil surface and 4-5 feet above the soil surface. Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off onto soil. Apply ONLY with a low volume sprayer operated at less than 20 PSI to prevent tree damage, bounce back and drift of spray droplets. Time applications to coincide with when most vulnerable pest life stage is present on plants. Do not apply to wet bark, during rainfall or if rain is expected within 12 hours. Control may be less effective in trees with thick bark, and at times when trees are not actively growing or transpiring. For Mountain Pine Beetle: apply from 2 weeks before to 2 weeks after expected peak of adult flight activity.

(continued)

ORNAMENTAL PLANTS AND FORESTS (continued)

BASAL TRUNK SPRAYS IN TREES AND LARGE SHRUBS (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Shrubs Ornamental Trees* Non-Bearing Fruit Trees Non-Bearing Nut Trees Trees in Plantations including: Conifer Deciduous Reforestation Nurseries Forests and Wooded Areas: National, Private and State	(continued) Pine Tip Moth (larvae) Psyllids Roundheaded Borers (excluding Asian Longhorned) Scales including: Calico Cryptomeria Elongate Hemlock Fig (Ficus) Wax Thrips (suppression) Walnut Twig Beetle Whiteflies including: Fig (Ficus)	12-24 oz per gallon Depending on bark type and thickness, one gallon of spray solution will typically cover 65-85" of cumulative trunk diameter (1.5-2.0 fl oz per inch of trunk diameter) when applied to trunk between soil surface and 4-5 feet above soil surface.	When sprayed on the trunk, <i>Safari 20 SG</i> Insecticide will be absorbed through the bark and into the vascular system, and then transported throughout the tree. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively transpiring trees, control may be evident within 1-3 weeks after application. Spray bark on root flare (buttress roots) and on trunk between soil surface and 4-5 feet above the soil surface. Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off onto soil. Apply ONLY with a low volume sprayer operated at less than 20 PSI to prevent tree damage, bounce back and drift of spray droplets. Time applications to coincide with when most vulnerable pest life stage is present on plants. Do not apply to wet bark, during rainfall or if rain is expected within 12 hours. Control may be less effective in trees with thick bark, and at times when trees are not actively growing or transpiring. For Mountain Pine Beetle: apply from 2 weeks before to 2 weeks after expected peak of adult flight activity.

(continued)

ORNAMENTAL PLANTS AND FORESTS (continued)
BASAL TRUNK SPRAYS IN TREES AND LARGE SHRUBS (continued)

Crops	Pests	Product Rate (By Weight)	Remarks
Christmas Trees Ornamental Trees* with trunk diameter less than 3" at soil line	Elongate Hemlock Scale Cryptomeria Scale Ficus (Fig) Whitefly	1.5-6.0 oz/gallon One gallon of spray solution will typically cover 325-425" of cumulative trunk diameter (0.3-0.4 fl oz per inch of trunk diameter) when applied to trunk between soil surface and 1 foot above soil surface.	For Christmas trees and ornamental trees less than 3" in diameter at soil line, spray trunk just to point of runoff between soil surface and 12" above soil surface.

One (1) level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz by weight of *Safari* 20 SG Insecticide.

Restrictions:

Do not apply more than 2.7 lbs (0.54 lbs ai) per acre of nursery, forest or landscape per year.

* See species restrictions in the **Application to Ornamental Plants (including Forestry)** section under **Application Information**.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool dry place. Do not store diluted spray. For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-CLEANUP.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration.

CONDITIONS OF SALE

Valent U.S.A. LLC warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions to the crops specified. To the extent consistent with applicable law, there are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

©2019 Valent U.S.A. LLC

Safari is a registered trademark of Valent U.S.A. LLC

Manufactured for

Valent U.S.A. LLC

P.O. Box 8025

Walnut Creek CA 94596-8025

Made in U.S.A.

Form 1812-F

TAJ 15AUG16


EPA Reg. No. 86203-11-59639

EPA Est. 67545-AZ-1

GROUP 4A INSECTICIDE

Safari®

20 SG INSECTICIDE



**FOR FOLIAR AND SYSTEMIC INSECT CONTROL IN
ORNAMENTAL PLANTS AND VEGETABLE TRANSPLANTS
IN ENCLOSED STRUCTURES.**

**FOR GREENHOUSE, NURSERY, INTERIOR PLANTSCAPE,
OUTDOOR LANDSCAPE AND FORESTRY USE ONLY.**

Active Ingredient:
Dinotefuran, [N-methyl-N'-nitro-N''-
((tetrahydro-3-furanyl)methyl)guanidine]... 20%
Other Ingredients 80%
Total 100%

EPA Reg. No. 86203-11-59639
EPA Est. 67545-AZ-1

Valent U.S.A. LLC
P.O. Box 8025, Walnut Creek CA 94596-8025

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

**SEE INSIDE BOOKLET FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

NET CONTENTS 12 OUNCES Form 1812-F



Safety Data Sheet - GHS

1. IDENTIFICATION: CHEMICAL PRODUCT AND COMPANY

PRODUCT NAME: Safari™ 20 SG Insecticide
EPA REGISTRATION NUMBER: 86203-11-59639; 33657-16-59639
VC NUMBER(S): 1455
SYNONYM(S): Dinotefuran 20% SG
PRODUCT DESCRIPTION: Insecticide for greenhouse, nursery, interior plantscape and outdoor landscape use.

Safari is a registered trademark of Valent U.S.A. LLC

MANUFACTURER/DISTRIBUTOR
 VALENT U.S.A. LLC
 P.O. Box 5075
 4600 Norris Canyon Road
 San Ramon, CA 94583

EMERGENCY TELEPHONE NUMBERS
 HEALTH EMERGENCY OR SPILL (24 hr):
 (800) 892-0099
 TRANSPORTATION (24 hr.): CHEMTREC
 (800) 424-9300 or (202) 483-7616

PRODUCT INFORMATION
 PROFESSIONAL PRODUCTS: (800) 898-2536

The current SDS is available through our website (www.valent.com), or by calling the product information numbers listed above.

2. HAZARDS IDENTIFICATION

This product is an EPA FIFRA registered pesticide. Some classifications on this SDS are not the same as the FIFRA-required classifications on the product label. Certain sections of this SDS are superseded by federal law under EPA FIFRA for a registered pesticide. Please see Section 15, REGULATORY INFORMATION for an explanation.

Classification - (per U.S. OSHA 29 CFR 1910.1200 (Hazcom 2012))

Acute toxicity - Inhalation (Dusts/Mists)	Category 4
Skin corrosion/irritation	Category 3

Label elements

EMERGENCY OVERVIEW

WARNING



Hazard statements

Harmful if inhaled
Causes skin irritation

Precautionary statements**Prevention**

Avoid breathing dust/fume/gas/mist/vapors/spray
Use only outdoors or in a well-ventilated area
Avoid contact with eyes and skin

Response

Eyes None.

Skin Wash thoroughly with soap and water. Wash contaminated clothing before reuse.

Inhalation IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTER or doctor/physician if you feel unwell.

Ingestion None.

FIRE None.

Spill None.

Storage

None

Disposal

None

Hazards not otherwise classified (HNOC)**Other Information**

- Toxic to aquatic life with long lasting effects
- Harmful to aquatic life

For information on Transportation requirements, see Section 14.

3. COMPOSITION/INFORMATION ON INGREDIENTS

Chemical name	CAS No.	Weight-%	TRADE SECRET
Dinotefuran Technical (99% ai)	165252-70-0	20.2	
Sodium dodecylbenzene sulfonate	25155-30-0	1 - 5	
Others	(Various CAS#s)	Balance of formulation	

* The chemical name, CAS number and/or exact percentage have been withheld as a trade secret

Other ingredients, which may be maintained as trade secrets, are any substances other than an active ingredient contained in this product. Some of these may be hazardous, but their identities are withheld because they are considered trade secrets. The hazards associated with the other ingredients are addressed in this document. Specific information on other ingredients for the management of exposures, spills, or safety assessments can be obtained by a treating physician or nurse by calling **(800) 892-0099** at any time.

4. FIRST AID MEASURES

EMERGENCY NUMBER (800) 892-0099

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Emergency Telephone: (800) 892-0099
REVISION NUMBER: 2

SDS NO.: 0426
REVISION DATE: 08/21/2020

You may also contact **1-800-892-0099** for emergency medical treatment information.

EYE CONTACT:

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

SKIN CONTACT:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

INGESTION:

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by the poison control center or doctor. Do not give anything to an unconscious person.

INHALATION:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTES TO PHYSICIAN:

None

5. FIRE FIGHTING MEASURES

Flash point °C Not Applicable

Flash point °F

EXTINGUISHING MEDIA: Water fog, carbon dioxide, foam, dry chemical

FLAMMABLE LIMITS IN AIR - LOWER (%):

Not applicable

FLAMMABLE LIMITS IN AIR - UPPER (%):

Not applicable

NFPA RATING:

Health:	1
Flammability:	3
Reactivity:	1
Special:	None

(Least-0, Slight-1, Moderate-2, High-3, Extreme-4). These values are obtained using professional judgement. Values were not available in the guidelines or published evaluations prepared by the National Fire Protection Association, NFPA.

FIRE FIGHTING INSTRUCTIONS: Products of combustion from fires involving this material may be toxic. Avoid breathing smoke and mists. Avoid personnel and equipment contact with fallout and runoff. Minimize the amount of water used for fire fighting. Do not enter any enclosed area without full protective equipment, including self-contained breathing equipment. Contain and isolate runoff and debris for proper disposal. Decontaminate personal protective equipment and fire fighting equipment before reuse.

This material is not expected to burn or explode in normal conditions, but will burn violently if involved in a fire. Dinotefuran becomes self-reactive in high temperatures. Exposure to heat may promote violent decomposition.

HAZARDOUS DECOMPOSITION PRODUCTS: Normal combustion forms carbon dioxide, water vapor and may produce: oxides of nitrogen,.

6. ACCIDENTAL RELEASE MEASURES

Emergency Telephone: (800) 892-0099
REVISION NUMBER: 2

SDS NO.: 0426
REVISION DATE: 08/21/2020

VALENT EMERGENCY PHONE NUMBER: (800) 892-0099
CHEMTREC EMERGENCY PHONE NUMBER: (800) 424-9300
OBSERVE PRECAUTIONS IN SECTION 8: PERSONAL PROTECTION

Stop the source of the spill if safe to do so. Contain the spill to prevent further contamination of the soil, surface water, or ground water. For additional spill response information refer to the North American Emergency Response Guidebook.

UN/NA NUMBER: Not applicable **EMERGENCY RESPONSE GUIDEBOOK NO.:** Not applicable

FOR SPILLS ON LAND:

CONTAINMENT: Remove all sources of ignition. Ventilate area of leak or spill. Clean-up personnel may require protection from inhalation of dust. Avoid runoff into storm sewers or other bodies of water.

CLEANUP: Clean up spill immediately in a manner that does not disperse dust into the air and place in a chemical waste container. Wash area with soap and water. Pick up wash liquid with additional absorbent and place in a chemical waste container.

FOR SPILLS IN WATER:

CONTAINMENT: This material will disperse or dissolve in water. Stop the source of the release. Contain and isolate to prevent further release on to soil or into surface water.

CLEANUP: Clean up spill immediately. Absorb spill with inert material. Remove contaminated water for treatment or disposal.

7. HANDLING AND STORAGE

END USER MUST READ AND OBSERVE ALL PRECAUTIONS ON PRODUCT LABEL.

HANDLING:

Keep away from all possible sources of ignition (sparks or flame). Avoid high temperatures exceeding 150°C. Keep container closed. Use only with adequate ventilation.

To avoid fire or explosion, dissipate static electricity during transfer by grounding and bonding containers and equipment before transferring the material. Use explosion-proof electrical equipment. Take precautionary measures against static discharges.

STORAGE:

Keep pesticide in original container. Do not store or transport near food or feed. Do not contaminate food or feed. Do not put concentrate into food or drink containers. Do not dilute concentrate in food or drink containers. Store in a cool, dry place, out of direct sunlight. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store diluted spray.

8. EXPOSURE CONTROLS/PERSONAL PROTECTION

END USER MUST READ AND OBSERVE ALL PRECAUTIONS ON PRODUCT LABEL.

INFORMATION FOR END USERS

Mixers, loaders, applicators and other handlers should refer to the product label before use for detailed information on personal protective equipment (PPE).

EYES & FACE: Do not get this material in your eyes. Eye contact can be avoided by wearing protective eyewear.

RESPIRATORY PROTECTION: Use this material only in well ventilated areas. If operating conditions result in airborne concentrations of this material, the use of an approved respirator is recommended.

SKIN & HAND PROTECTION: Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks and chemical-resistant gloves made of any waterproof material.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS: Use in a well ventilated area.

EXPOSURE LIMITS

Chemical name	ACGIH Exposure Limits	OSHA Exposure Limits	Manufacturer's Exposure Limits
Dinotefuran Technical (99% ai)	None	None	None
Sodium dodecylbenzene sulfonate	None	None	None
Others	None	None	None

9. PHYSICAL AND CHEMICAL PROPERTIES

Physical state	Solid	Odor	Odorless
Appearance	Granules	Odor threshold	No information available
Color	Off-white		

<u>PROPERTIES</u>	<u>Values</u>	<u>Remarks • Method</u>
pH	7.6	(1% aqueous solution)
Melting point/freezing point	107.5 °C	Melting point (Dinotefuran Technical)
Boiling point/boiling range	Decomposed at 208°C (Technical)	
Flash point	Not Applicable	
Evaporation rate	No information available	
Flammability (solid, gas)	No information available	
Flammability Limits in Air		
Upper flammability limits	No information available	
Lower flammability limit	No information available	
Vapor pressure	No information available	
Vapor density	No information available	
Specific Gravity	No information available	
Water solubility	Soluble in water	
Solubility in other solvents	No information available	
Partition coefficient	No information available	
Autoignition temperature	No information available	
Decomposition temperature	No information available	
Viscosity	No information available	
Explosive properties	No information available	
Oxidizing properties	No information available	
Liquid Density	No information available	
Bulk density	0.56 g/mL	

10. STABILITY AND REACTIVITY

Reactivity

Not an oxidizing or reducing agent.

Chemical stability

Stable under normal ambient conditions.

Possibility of Hazardous Reactions

This material is combustible and may form explosive dust-air mixture.

Conditions to avoid

Extremes of temperature and direct sunlight.

Incompatible materials

Heat and ignition sources. Oxidizers.

Hazardous Decomposition Products

Normal combustion forms carbon dioxide, water vapor and may produce: oxides of nitrogen.

11. TOXICOLOGICAL INFORMATION

ACUTE TOXICITY:

Oral Toxicity LD ₅₀ (rats)	> 2,000 mg/kg	EPA Tox Category	III
Dermal Toxicity LD ₅₀ (rabbits)	> 2,000 mg/kg	EPA Tox Category	III
Inhalation Toxicity LC ₅₀ (rats)	> 2.94 mg/L (4 hr)	EPA Tox Category	IV
Eye Irritation (rabbits)	Moderately irritating	EPA Tox Category	III
Skin Irritation (rabbits)	Mildly irritating	EPA Tox Category	III
Skin Sensitization (guinea pigs)	Non-sensitizer	EPA Tox Category	Not applicable

CARCINOGEN CLASSIFICATION

Chemical name	IARC Group 1 or 2	OSHA - Select Carcinogens	NTP Carcinogen List
Others	Not listed	Not listed	Not listed
Dinotefuran Technical (99% ai)	Not listed	Not listed	Not listed
Sodium dodecylbenzene sulfonate	Not listed	Not listed	Not listed

TOXICITY OF DINOTEFURAN TECHNICAL

SUBCHRONIC: Dinotefuran technical was tested in 13-week dietary toxicity studies in rats, mice and dogs. In the rat study, a NOEL of 500 ppm was established, based on reduced body weight gain in females and adrenal cortical vacuolation in males and a NOAEL of 5,000 ppm based on marked growth retardation at 25,000 ppm (adrenal cortical vacuolation not adverse). A NOEL of 25,000 ppm was established in the mouse study based on reduced body weight gain at 50,000 ppm. In the dog 13-week dietary study, a NOEL of 8,000 ppm was established based on reduced body weight gain. No target organs were identified in subchronic inhalation or dermal toxicity studies in rats.

CHRONIC/CARCINOGENICITY: Dinotefuran technical was tested in lifetime studies with rats and mice and a one-year study with dogs. In common with the subchronic studies in these species, no specific target organs could be identified. In the 78-week mouse study a NOAEL of 2500 ppm was established, based on decreased weight gain and a decrease in circulating platelet counts. In the 104-week rat study a NOAEL of 2000 ppm was established, based on a decrease in weight gain in females. There were no treatment-related effects in rats or mice on survival or the nature and incidence of neoplastic and adverse non-neoplastic histomorphological findings in either species at any dose level. In the 52-week dog study a NOAEL of 16000 ppm was established based on decreased weight gain in both sexes and decreased food consumption in females.

NEUROTOXICITY: Dinotefuran did not produce any functional or histomorphological evidence of neurotoxicity in acute (gavage) and 13-week (dietary) neurotoxicity studies in rats. The NOEL for neurotoxicity in the acute study was 1,500 mg/kg, the highest dose level administered. The NOEL for neurotoxicity in the 13-week dietary study was

50,000 ppm. The NOEL for all effects in this study was 5,000 ppm based on reduced body weight gain and food consumption.

DEVELOPMENTAL TOXICITY: In a developmental toxicity study of Dinotefuran technical in rats the maternal NOEL was 300 mg/kg/day based on reduced weight gain, food consumption and water intake at 1000 mg/kg/day. Dinotefuran technical did not produce developmental effects in rats at doses up to 1000 mg/kg/day (the highest does tested). In a study with rabbits the maternal NOEL was 52 mg/kg/day based on reduced weight gain, food consumption and water intake and clinical signs noted at 300 mg/kg/day and pathology findings in the liver and stomach at 125 mg/kg/day and higher. The developmental NOEL was 300 mg/kg/day.

REPRODUCTION: Dinotefuran technical was tested in a two-generation rat reproduction study at doses of 0, 300, 1000, 3000 and 10000 ppm. The NOEL for systemic toxicity in parental animals was 3000 ppm based on decreased body weight gain and food consumption and decreased spleen and thyroid weights at the highest dose level evaluated (10000 ppm). The NOEL for reproductive effects was 10000 ppm. The NOEL for effects on the offspring was 3000 ppm based on reduced preweaning weight gain at 10000 ppm.

MUTAGENICITY: Dinotefuran technical was negative in the following in vitro assays: Ames Assay, mouse lymphoma (L5178Y), mammalian cytogenetics (CHL/IU) or DNA Repair. Dinotefuran technical was negative in the following in vivo assays: mouse micronucleus. Overall, Dinotefuran technical does not present a genetic hazard.

For a summary of the potential for adverse health effects from exposure to this product, refer to Section 2. For information regarding regulations pertaining to this product, refer to Section 15.

12. ECOLOGICAL INFORMATION

AVIAN TOXICITY: Dinotefuran Technical is practically non-toxic to moderately toxic to avian species. Test results include:
 Oral LD₅₀ quail: greater than 2000 mg/kg
 Dietary LC₅₀ Mallard duck: greater than 997.9 ppm
 Dietary LC₅₀ quail: greater than 1301 ppm
 Reproduction quail: NOEL = 5000 ppm
 Reproduction Mallard duck: NOEL = 2000 ppm

AQUATIC ORGANISM TOXICITY: Dinotefuran Technical is practically non-toxic to fish and ranges from practically nontoxic to highly toxic to aquatic invertebrate species (especially shrimp.) Test results include:
 LC₅₀ (96 hr) Bluegill Sunfish: greater than 100 mg/L
 LC₅₀ (96 hr) Rainbow Trout: greater than 100 mg/L
 LC₅₀ (96 hr) Common Carp: greater than 100 mg/L
 LC₅₀ (96 hr) Sheepshead Minnow: greater than 109 mg/L
 NOEC (early life stage) Rainbow Trout: greater than 10 mg/L
 EC₅₀ (48 hr) Daphnia magna: greater than 1000 mg/L
 NOEC (lifecycle) Daphnia magna: > 10 mg/L
 LC₅₀ (96 hr) saltwater Mysid Shrimp: 0.79 mg/L
 NOEC (lifecycle) saltwater Mysid Shrimp: 0.089mg/L
 EC₅₀ (96 hr) Oyster Shell Deposition: greater than 141 mg/L
 ErC₅₀ (0-72 hr) Algae (P. subcapitata): greater than 100 mg/L

OTHER NON-TARGET ORGANISM TOXICITY:

Dinotefuran Technical is highly toxic to bees. The acute oral and contact LD₅₀ in bees were 0.056 µg/bee and 0.022 ug/bee, respectively. This product is highly toxic to bees or other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees or other pollinating insects are foraging in the treatment

area.

OTHER ENVIRONMENTAL INFORMATION:

This pesticide is toxic to shrimp. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below mean high water mark. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

13. DISPOSAL CONSIDERATIONS

END USERS MUST DISPOSE OF ANY UNUSED PRODUCT AS PER THE LABEL RECOMMENDATIONS.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Completely empty container by shaking and tapping sides and bottom to loosen clinging particles promptly after emptying. Then, offer for recycling if available, or reconditioning if appropriate, or puncture and dispose of container in a sanitary landfill, or by other procedures approved by state and local authorities. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration.

DISPOSAL METHODS: Check government regulations and local authorities for approved disposal of this material. Dispose of in accordance with applicable laws and regulations.

14. TRANSPORTATION INFORMATION

DOT (ground) SHIPPING NAME: Not regulated for domestic ground transport by U.S. DOT
REMARKS: None
EMERGENCY RESPONSE Not applicable
GUIDEBOOK NO.:

ICAO/IATA SHIPPING NAME: UN 3077 Environmentally Hazardous Substance, Solid, N.O.S. (Dinotefuran), 9, III, Marine Pollutant
REMARKS:
 •Single or inner packaging less than 5 L (liquid) or 5 Kg net (solids) excepted from Dangerous Goods regulations – see IATA Special Provision A197.
 •For US shipping, Emergency Response Guidebook No. 171

IMDG SHIPPING NAME: UN 3077 Environmentally Hazardous Substance, Solid, N.O.S. (Dinotefuran), 9, III, Marine Pollutant
REMARKS:
 •Single or inner packaging less than 5 L (liquid) or 5 Kg net (solids) excepted from Dangerous Goods regulations – see IMDG 2.10.2.7
 •For US shipping, Emergency Response Guidebook No. 171
EMS NO.: F-A, S-F

15. REGULATORY INFORMATION

EPA-FIFRA LABEL INFORMATION THAT DIFFERS FROM OSHA-GHS REQUIREMENTS:

Pesticide products in the U.S. are registered by the EPA under FIFRA and are subject to certain labeling requirements under federal pesticide law. These requirements may differ from the classification criteria and hazard information

required by OSHA GHS for safety data sheets, and for workplace labels of non-pesticide chemicals. The following is the hazard information as required on the FIFRA pesticide label:

EPA FIFRA SIGNAL WORD: CAUTION

- *Harmful if swallowed*
- *Harmful if absorbed through skin*
- *Causes moderate eye irritation*
- *Avoid contact with eyes, skin and clothing.*
- *Wash hands thoroughly with soap and water after handling.*
- *Remove contaminated clothing and wash before re-use.*
- *Keep out of reach of children.*

PESTICIDE REGULATIONS: All pesticides are governed under FIFRA (Federal Insecticide, Fungicide, and Rodenticide Act). Therefore, the regulations presented below are pertinent only when handled outside of the normal use and applications of pesticides. This includes waste streams resulting from manufacturing/formulation facilities, spills or misuse of products, and storage of large quantities of products containing hazardous or extremely hazardous substances.

U.S. FEDERAL REGULATIONS: Ingredients in this product are reviewed against an inclusive list of federal regulations. Therefore, the user should consult appropriate authorities. The federal regulations reviewed include: Clean Water Act, SARA, CERCLA, RCRA, DOT, TSCA and OSHA. If no components or information is listed in the space below this paragraph, then none of the regulations reviewed are applicable.

Sodium dodecylbenzene sulfonate

Clean Water Act - Hazardous Substances	Present
CERCLA Reportable Quantity (RQ):	1000 lb 454 kg

SARA (311, 312):

Immediate Health:	Yes
Chronic Health:	No
Fire:	Yes
Sudden Pressure:	No
Reactivity:	No

STATE REGULATIONS: Each state may promulgate standards more stringent than the federal government. This section cannot encompass an inclusive list of all state regulations. Therefore, the user should consult state or local authorities. The state regulations reviewed include: California Proposition 65, California Directors List of Hazardous Substances, Massachusetts Right to Know, Michigan Critical Materials List, New Jersey Right to Know, Pennsylvania Right to Know, Rhode Island Right to Know and the Minnesota Hazardous Substance list. For Washington State Right to Know, see Section 8 for Exposure Limit information. For Louisiana Right to Know refer to SARA information listed under U.S. Regulations above. If no components or information is listed in the space below this paragraph, then none of the regulations reviewed are applicable.

Sodium dodecylbenzene sulfonate

California - Directors List of Hazardous Substances	Present
MA Right To Know	Present
NJ Right To Know	1698
PA Right To Know	Environmental hazard

For information regarding potential adverse health effects from exposure to this product, refer to Sections 2 and 11.

16. OTHER INFORMATION

REASON FOR ISSUE: Updated Manufacturer/Distributor Address and General Review.
SDS NO.: 0426
EPA REGISTRATION NUMBER: 86203-11-59639; 33657-16-59639
REVISION NUMBER: 2
REVISION DATE: 08/21/2020
SUPERCEDES DATE: 08/21/2020
RESPONSIBLE PERSON(S): Valent U.S.A. LLC, Corporate EH&S, (925) 256-2803

The information provided in this Safety Data Sheet (SDS) is provided in good faith and believed to be accurate at the time of preparation of the SDS. However, to the extent consistent with applicable law, Valent U.S.A. LLC and its subsidiaries or affiliates extend no warranties, make no representations, and assume no responsibility as to the accuracy, suitability, or completeness of such information. Additionally, to the extent consistent with applicable law, neither Valent U.S.A. LLC nor any of its subsidiaries or affiliates represents or guarantees that this information or product may be used without infringing the intellectual property rights of others. Except to the extent a particular use and particular information are expressly stated on the product label, it is the users' own responsibility to determine the suitability of this information for their own particular use of this product. If necessary, contact Valent U.S.A. LLC to confirm that you have the most current product label and SDS.

This SDS provides important health, safety, and environmental information for employers, employees, emergency responders and others handling large quantities of the product in activities generally other than product use as required by the Occupational Health and Safety Act (29 CFR 1910.1200, "Hazcom").

The product label provides information specifically for product use in the ordinary course. All necessary hazard classification and appropriate precautionary use, storage, and disposal information is set forth on that label or labeling accompanying the pesticide or to which reference is made on the label.

2020 Valent U.S.A. LLC